

Final Report

Ground Truthing of Exotic Species Monitoring on District-Managed Lands in South Florida

Submitted by:

Keith Bradley, Steven Woodmansee, and George Gann
The Institute for Regional Conservation
22601 SW 152 Ave.
Miami, Florida 33170

To

The South Florida Water Management District
3301 Gun Club Road
West Palm Beach, Florida 33406

November 10, 1997

I. Introduction

From June through October 1997 The Institute for Regional Conservation conducted botanical surveys of seven properties managed by the South Florida Water Management District. The purpose of these surveys was to determine the status of non-native exotic taxa on these lands, as well as the status of endangered and threatened taxa. These properties include The Frog Pond, Loxahatchee River Corridor, Pal-Mar (West Jupiter Wetlands), Water Conservation Area 2, Nicodemus Slough, Six Mile Cypress, and the marshes along the southwest end of Lake Okeechobee (from Fisheating Creek to Torry & Kreamer Islands).

II. Methods

Surveys were conducted by one field researcher and one technician. These surveys were made at intervals from June through October in order to detect seasonally flowering, ephemeral taxa (note that some taxa which only bloom in the winter and spring may not have been detected in these surveys). All vascular plants seen within each property were recorded. For all taxa listed as exotic by Wunderlin, et. al. (1996, Atlas of Florida Vascular Plants) estimates of population size, acreage, and basic habitat were made. The same was done for all taxa listed by the U.S. Fish and Wildlife Service or Florida Division of Plant Industry as Endangered or Threatened. The population status (increasing, decreasing, stable) and the number of populations was also estimated for each of these listed taxa. Attempts were made to collect herbarium voucher specimens of each of these taxa, as long as they were of sufficient quality to be placed in a museum collection (e.g. flowering and/or fruiting, without insect damage).

III. Results

Attached are tables of the plant taxa recorded at each site. Note that species in bold type are considered to be exotic, and species in italics are only present as cultivated, non-naturalized individuals. Taxonomy generally follows Wunderlin, et. al. (1996) Summary data included on each list for exotic plant taxa is as follows:

Nat. Area?: Yes if the taxon is invasive on relatively intact natural areas on the site, No if the taxon is restricted to disturbed areas.

Habitat: Soil moisture condition in which the taxon is found most frequently (mesic, xeric, or wet)

Acreage: Approximate area covered by the taxon based on a \log_{10} scale.

Pop. Size: Approximate number of individuals on the site based on a \log_{10} scale.

Florida Range: General range of the plant in Florida. Four categories are recognized: 1) *South* – including Martin, Glades, and Charlotte counties and south; 2) *Central* – north to include Volusia, Marion, and Levy counties; 3) *North* – north to the Georgia border and west to the Suwanee River, to include Gilchrist, Suwanee, and Columbia counties; and 4) *Panhandle*, west of the Suwanee River.

Natural Range: Natural range of the taxon.

Voucher: Number of herbarium voucher collection. BW = Bradley and Woodmansee, GB = Gann and Bradley, B = Bradley.

Also attached are tables listing endangered and threatened taxa found on each site. Summary data included on each list is as follows:

State: Status of each taxon on the Official List of Threatened and Endangered Species, Florida Department of Agriculture and Consumer Services.

Habitat: Soil moisture condition in which the taxon is found most frequently (mesic, xeric, or wet).

Acreage: Approximate area covered by the taxon based on a \log_{10} scale.

Pop. Size: Approximate number of individuals on the site based on a \log_{10} scale.

Status: Estimate of population trend (decreasing, increasing, or stable).

Stations: Approximate number of populations on the site. Some taxa are widespread on the site and these are noted by the designation "Throughout."

Native Range: Natural Range of the taxon.

Voucher: Number of herbarium voucher collection. BW = Bradley and Woodmansee, GB = Gann and Bradley, B = Bradley.

Exotic Taxa

A total of 188 exotic plant taxa were recorded. Of these 63 (33.5%) are considered to be invasive on one or more of these sites. Following is a summary for each site:

| | #Taxa ¹ | #Exotics ² | #Invasive ³ | %Exotic | % Invasive ⁴ |
|----------------------------|--------------------|-----------------------|------------------------|---------|-------------------------|
| Frog Pond | 395 | 87 | 14 | 22.0 | 16.1 |
| Lake Okeechobee Marshes | 187 | 43 | 14 | 23.0 | 32.6 |
| Loxahatchee River Corridor | 574 | 112 | 43 | 19.5 | 38.4 |
| Nicodemus Slough | 256 | 33 | 14 | 12.9 | 42.4 |
| Pal-Mar | 350 | 35 | 10 | 10.0 | 28.6 |
| Six Mile Cypress | 431 | 48 | 24 | 11.1 | 50.0 |
| Water Conservation Area 2 | 272 | 54 | 12 | 19.9 | 22.2 |

See Table I "Exotic Plant Taxa" for a list of all exotic plant taxa recorded during these surveys, and the number of sites each taxon was recorded at. See Table II "Invasive Taxa" for a list of all invasive taxa, the frequency at which they were found at the seven properties, the total number of sites each taxon was recorded at, and the percentage of sites each taxon was invasive at.

Endangered and Threatened Taxa

A total of 20 endangered and 8 threatened plant taxa were recorded. Following is a summary for each site:

| | #Endangered | #Threatened | Total |
|----------------------------|-------------|-------------|-------|
| Frog Pond | 9 | 4 | 13 |
| Lake Okeechobee Marshes | 2 | 0 | 2 |
| Loxahatchee River Corridor | 7 | 5 | 12 |
| Nicodemus Slough | 1 | 2 | 3 |
| Pal-Mar | 4 | 4 | 8 |
| Six Mile Cypress | 6 | 5 | 11 |
| Water Conservation Area 2 | 0 | 0 | 0 |

See Table III "Endangered and Threatened Taxa" for a list of endangered and threatened taxa and the frequency at which they were found on the seven properties.

¹ Excluding cultivated taxa

² Excluding cultivated taxa

³ Taxa which are recruiting in relatively undisturbed natural areas

⁴ % of Exotic taxa which are invasive

Frog Pond - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| Voucher | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range |
|---|------------|---------|-----------|----------------|-----------------------|------------------|
| <i>Acacia farnesiana</i> | No | Mesic | 101-1,000 | 1,001-10,000 | South, Central | Tropical America |
| <i>Acalypha alopecuroides</i> | | | | | | B 1225 |
| <i>Acalypha ostryifolia</i> | | | | | | |
| <i>Acrostichum danaeifolium</i> | | | | | | |
| <i>Aeschynomene americana</i> | | | | | | |
| <i>Agalinis fasciculata</i> | | | | | | |
| <i>Agalinis linifolia</i> | | | | | | |
| <i>Aletis lutea</i> | Yes | Wet | 1-100 | 1,001-10,000 | Throughout | South America |
| <i>Alternanthera philoxeroides</i> | | | | | | BW 354 |
| <i>Anmannia coccinea</i> | | | | | | |
| <i>Ammannia latifolia</i> | | | | | | |
| <i>Amaranthus spinosus</i> | No | Mesic | 101-1,000 | 10,001-100,000 | Throughout | India |
| <i>Amaranthus viridis</i> | No | Mesic | 101-1,000 | 10,001-100,000 | Throughout | Tropical America |
| <i>Ambrosia artemisiifolia</i> | | | | | | |
| <i>Ampelopsis arborea</i> | | | | | | |
| <i>Anagallis arvensis</i> | No | Mesic | <1 | 101-1,000 | Throughout | Eurasia |
| <i>Andropogon glomeratus</i> var. <i>pumilus</i> | | | | | | BW 2 |
| <i>Andropogon longiberbis</i> | | | | | | |
| <i>Andropogon ternarius</i> | | | | | | |
| <i>Anemia adiantifolia</i> | | | | | | |
| <i>Anemia wrightii</i> | | | | | | |
| <i>Angadenia berteroii</i> | | | | | | |
| <i>Ardisia elliptica</i> | No | Mesic | <1 | 101-1,000 | South, Central | India |
| <i>Ardisia esculentoides</i> | | | | | | |
| <i>Aristida purpurascens</i> var. <i>tenuispica</i> | No | Mesic | <1 | 101-1,000 | South, Central | South America |
| <i>Asclepias curassavica</i> | | | | | | BW 346 |
| <i>Asclepias lanceolata</i> | | | | | | |
| <i>Aster adnatus</i> | | | | | | |
| <i>Aster bracteatus</i> | | | | | | |
| <i>Aster dumosus</i> | | | | | | |
| <i>Aster subulatus</i> | | | | | | |
| <i>Axonopus furcatus</i> | | | | | | |
| <i>Azolla caroliniana</i> | | | | | | |
| <i>Baccharis glomeruliflora</i> | | | | | | |
| <i>Baccharis halimifolia</i> | | | | | | |
| <i>Bacopa caroliniana</i> | | | | | | |
| <i>Bacopa monnieri</i> | | | | | | |
| <i>Bidens alba</i> var. <i>radiata</i> | | | | | | |
| <i>Bischofia javanica</i> | No | Mesic | <1 | 101-1,000 | South, Central | Tropical Asia |
| <i>Blechum pyramidatum</i> | No | Mesic | <1 | 1-100 | South, Central, North | Tropical America |
| <i>Bletia purpurea</i> | | | | | | |
| <i>Boehmeria cylindrica</i> | | | | | | |
| <i>Bothriochloa pertusa</i> | No | Mesic | <1 | 101-1,000 | South, Central | Old World |
| <i>Brassica juncea</i> | No | Mesic | <1 | 1-100 | Throughout | Europe |
| <i>Buchnera americana</i> | | | | | | B 1180 |

Frog Pond - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| Voucher | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range |
|--|------------|---------|-----------|--------------|-----------------------|-------------------------------------|
| <i>Bucida buceras</i> | | | | | | |
| <i>Bursera simaruba</i> | | | | | | |
| <i>Byrsinima lucida</i> | | | | | | |
| <i>Callicarpa americana</i> | | | | | | |
| <i>Callistemon viminalis</i> | | | | | | |
| <i>Caperonia palustris</i> | No | Wet | <1 | 101-1,000 | South, Central | Tropical America |
| <i>Capraria biflora</i> | | | | | | |
| <i>Cardamine pensylvanica</i> | | | | | | |
| <i>Cardiospermum halicacabum</i> | No | Mesic | <1 | 1-100 | South, Central | Tropical America |
| <i>Carica papaya</i> | Yes | Mesic | 1-100 | 1,001-10,000 | South, Central | Tropical America |
| <i>Cassytha filiformis</i> | | | | | | |
| <i>Casuarina equisetifolia</i> | Yes | Mesic | 101-1,000 | 1,001-10,000 | South, Central, North | Australia |
| <i>Cenchrus echinatus</i> | | | | | | |
| <i>Centella asiatica</i> | | | | | | |
| <i>Ceratopteris thalictroides</i> | | | | | | |
| <i>Chamaecrista nictitans</i> var. <i>aspera</i> | | | | | | |
| <i>Chamaesyce blodgettii</i> | | | | | | |
| <i>Chamaesyce conferta</i> | | | | | | |
| <i>Chamaesyce hirta</i> | | | | | | |
| <i>Chamaesyce hypericifolia</i> | | | | | | |
| <i>Chamaesyce hyssopifolia</i> | No | Mesic | <1 | 101-1,000 | South | ? |
| <i>Chamaesyce lastiocarpa</i> | | | | | | |
| <i>Chamaesyce mendezii</i> | | | | | | |
| <i>Chamaesyce ophthalmica</i> | | | | | | |
| <i>Chamaesyce prostrata</i> | | | | | | |
| <i>Chaptalia albicans</i> | | | | | | |
| <i>Chenopodium album</i> | No | Mesic | <1 | 101-1,000 | Throughout | Europe |
| <i>Chenopodium murale</i> | No | Mesic | <1 | 101-1,000 | South, Central, North | Europe |
| <i>Chiococca alba</i> | | | | | | |
| <i>Chiococca parvifolia</i> | | | | | | |
| <i>Chloris cf. radiata</i> | No | Mesic | <1 | 1-100 | South | Tropical America |
| <i>Chromolaena odorata</i> | | | | | | |
| <i>Cirsium horridulum</i> | | | | | | |
| <i>Cissus verticillata</i> | | | | | | |
| <i>Citharexylum spinosum</i> | | | | | | |
| <i>Cladium jamaicense</i> | | | | | | |
| <i>Coccocypselma diversifolia</i> | | | | | | |
| <i>Coelorrhiza rugosa</i> | No | Wet | 1-100 | 1,001-10,000 | Throughout | Asia |
| <i>Colocasia esculenta</i> | No | Wet | <1 | 101-1,000 | South, Central, North | Old World |
| <i>Commelinaceae</i> | | | | | | |
| <i>Commelinopsis diffusa</i> | | | | | | |
| <i>Conobea multifida</i> | No | Mesic | 101-1,000 | 1,001-10,000 | South | Georgia to Texas, Kansas, and Ohio. |
| <i>Conocarpus erectus</i> | | | | | | |
| <i>Conoclinium coelestinum</i> | | | | | | |
| <i>Conyza canadensis</i> var. <i>pusilla</i> | | | | | | |
| <i>Corchorus aestuans</i> | No | Mesic | 101-1,000 | 1,001-10,000 | Throughout | Asia |

Frog Pond - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|-----------|----------------|---------------|--|----------------|
| Corchorus siliquosus | | | | | | | |
| Coronopus diethylmus | No | Mesic | <1 | 101-1,000 | Throughout | Unknown, South America or Eurasia | BW 3, B 1187 |
| Crinum americanum | | | | | | | |
| Crotalaria pumila | | | | | | | |
| Cucumis melo | No | Mesic | <1 | 1-100 | Throughout | Old World Tropics | BW 431 |
| Cynanchum blodgettii | | | | | | | |
| Cynanchum scoparium | | | | | | | |
| Gnidodon dactylon | No | Mesic | 101-1,000 | >100,000 | Throughout | Africa | B 1208 |
| Cyperus compressus | | | | | | | |
| Cyperus croceus | | | | | | | |
| Cyperus elegans | | | | | | | |
| Cyperus esculentus | No | Mesic | 101-1,000 | >100,000 | Throughout | Old World | BW 361 |
| Cyperus flavescens | | | | | | | |
| Cyperus hastian | | | | | | | |
| Cyperus lata | No | Mesic | 101-1,000 | >100,000 | Throughout | Eastern Asia | BW 362 |
| Cyperus ligularis | | | | | | | |
| Cyperus ochraceus | No | Mesic | 1-100 | 1,001-10,000 | South | Coastal Plain, outside of Florida, West Indies, Mexico, central, America | B 1181 |
| Cyperus odoratus | | | | | | | |
| Cyperus polystachyos | | | | | | | |
| Cyperus rotundus | No | Mesic | 101-1,000 | >100,000 | Throughout | Eurasia | BW 370, B 1220 |
| Cyperus strigosus | | | | | | | |
| Cyperus surinamensis | | | | | | | |
| Dactyloctenium aegyptium | No | Mesic | 1-100 | 1,001-10,000 | Throughout | Old World Tropics | B 1223 |
| Desmodium regia | No | Mesic | <1 | 1-100 | South | Madagascar | |
| Desmodanthus cf. leptophyllus | No | Mesic | <1 | 1-100 | South | ? | |
| Desmodium tortuosum | | | | | | | |
| Dichanthelium commutatum | | | | | | | |
| Dichanthelium dichotomum | | | | | | | |
| Dichanthelium erectifolium | | | | | | | |
| Digitaria ciliaris | | | | | | | |
| Digitaria virginiana | | | | | | | |
| Diodia angustifolia | | | | | | | |
| Dipteris latifolia | | | | | | | |
| Dyschoriste angusta | | | | | | | |
| Echinocloa colonia | No | Mesic | 101-1,000 | 10,001-100,000 | Throughout | Old World Tropics | BW 368 |
| Echinocloa crusgalli | No | Mesic | <1 | 101-1,000 | Throughout | Eurasia | GB 1029 |
| Echinochloa walteri | | | | | | | |
| Eclipta prostrata | | | | | | | |
| Eleocharis cellulosa | | | | | | | |
| Eleocharis geniculata | No | Mesic | 1-100 | 1,001-10,000 | Throughout | Old World | |
| Eruca sativa | | | | | | | |
| Eruca sativa var. angustifolia | | | | | | | |
| Erythrina caroliniensis | | | | | | | |
| Erythrina caroliniensis var. angustifolia | | | | | | | |
| Erigeron canadensis | | | | | | | |
| Eragrostis ciliaris | No | Mesic | <1 | 1,001-10,000 | Throughout | West Indies to South America | B 1224 |
| Eragrostis elliotii | | | | | | | |
| Erechtites hieracifolia | | | | | | | |

Frog Pond - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---------------------------------|------------|---------|---------|--------------|----------------|--|---------|
| Eragrostis sp. | No | Mesic | <1 | 101-1,000 | Throughout | Southeast Asia | B 1188 |
| Eremochloa ophiurooides | No | Mesic | <1 | 1,001-10,000 | | | |
| <i>Eriogonum quercifolius</i> | | | | | | | |
| <i>Eugenia axillaris</i> | | | | | | | |
| <i>Eulophia alta</i> | | | | | | | |
| <i>Eupatorium capillifolium</i> | | | | | | | |
| <i>Eupatorium leptophyllum</i> | | | | | | | |
| <i>Eupatorium mikanioides</i> | | | | | | | |
| <i>Eupatorium serotinum</i> | | | | | | | |
| <i>Eustachys glauca</i> | | | | | | | |
| <i>Eustachys petraea</i> | | | | | | | |
| <i>Eustoma exaltata</i> | | | | | | | |
| <i>Evolvulus sericeus</i> | | | | | | | |
| <i>Ficus aurea</i> | | | | | | | |
| <i>Ficus citrifolia</i> | | | | | | | |
| <i>Fimbristylis cymosa</i> | | | | | | | |
| <i>Flaveria lineans</i> | | | | | | | |
| <i>Flaveria trinervia</i> | | | | | | | |
| <i>Flirena breviseta</i> | | | | | | | |
| <i>Galactia volubilis</i> | | | | | | | |
| Gallinsoga quadriradiata | No | Mesic | <1 | 1,001-10,000 | Throughout | Tropical America | BW 369 |
| <i>Galium hispidulum</i> | | | | | | | |
| <i>Galium tritectorium</i> | | | | | | | |
| <i>Gaura angustifolia</i> | | | | | | | |
| <i>Geranium carolinianum</i> | | | | | | | |
| <i>Guapira discolor</i> | | | | | | | |
| <i>Guettarda scabra</i> | | | | | | | |
| <i>Habenaria floribunda</i> | | | | | | | |
| <i>Heliotropium polypyllum</i> | | | | | | | |
| <i>Herissantia crispa</i> | | | | | | | |
| Heteranthera limosa | No | Wet | <1 | 1-100 | South, Central | N.M., Colo., and Va., W.I., Mex., C.A., S.A. | BW 353 |
| <i>Hypericum hypericoides</i> | | | | | | | |
| <i>Hypoxis alata</i> | | | | | | | |
| <i>Ilex cassine</i> | | | | | | | |
| <i>Ilex krigiana</i> | | | | | | | |
| Imperata cylindrica | No | Wet | 1-100 | 101-1,000 | Throughout | Old World | B 1213 |
| <i>Ipomoea heterophaea</i> | | | | | | | |
| <i>Ipomoea hederifolia</i> | | | | | | | |
| <i>Ipomoea sagittata</i> | | | | | | | |
| <i>Ipomoea cordatotriloba</i> | | | | | | | |
| <i>Jacquemontia curtissii</i> | | | | | | | |
| <i>Jacquemontia pentanthos</i> | | | | | | | |
| <i>Juncus marginatus</i> | | | | | | | |
| <i>Juncus megacephalus</i> | | | | | | | |
| <i>Koanophyllum villosum</i> | | | | | | | |
| <i>Kosteletzkya virginica</i> | | | | | | | |

Frog Pond - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|-----------|----------------|---------------------------|-------------------|---------|
| Lantana camara | No | Mesic | 101-1,000 | 1,001-10,000 | Throughout | Tropical America | |
| Lantana involucrata | | | | | | | |
| Leersia hexandra | | | | | | | |
| Lemna valvidiana | | | | | | | |
| Lepidium virginicum | | | | | | | |
| Leptochloa fascicularis | | | | | | | |
| Leptochloa mucronata | No | Mesic | <1 | 101-1,000 | South, Central, North | Caribbean? | |
| Leucaena leucocephala | | | | | | | |
| Leucospora multifida | | | | | | | |
| Lindernia crustacea | No | Mesic | <1 | 10,001-100,000 | Throughout | Indo-Malaya | BW 356 |
| Lobelia glandulosa | | | | | | | |
| Ludwigia octovalvis | | | | | | | |
| Ludwigia alata | | | | | | | |
| Ludwigia decurrens | | | | | | | |
| Ludwigia repens | | | | | | | |
| Lytthrum alatum var. lanceolatum | No | Mesic | 101-1,000 | 10,001-100,000 | South, Central, Panhandle | Tropical America | BW 352 |
| Macrorhynchium lathyroides | | | | | | | |
| Magnolia virginiana | | | | | | | |
| Malvastrum corymbosum | No | Mesic | <1 | 1-100 | South, Central, Panhandle | Tropical America | B 1184 |
| Malvastrum coronandellianum | | | | | | | |
| Mecardonia acuminata ssp. peninsulae | | | | | | | |
| Mecardonia procumbens | | | | | | | |
| Melanthera angustifolia | | | | | | | |
| Melanthera paniculata | | | | | | | |
| Melilotus albus | No | Mesic | 1-100 | 1,001-10,000 | Panhandle | Europe | |
| Melochia cf. corchorifolia | No | Mesic | <1 | 1-100 | Throughout | Old World Tropics | B 1228 |
| Melothria pendula | | | | | | | |
| Merremia tuberosa | No | Mesic | <1 | 1-100 | South | Tropical America | |
| Melopodium toxiferum | | | | | | | |
| Mikania cordifolia | | | | | | | |
| Mikania scandens | | | | | | | |
| Mitreola petiolata | | | | | | | |
| Mitreola sessilifolia | | | | | | | |
| Momordica charantia | No | Mesic | 101-1,000 | 1,001-10,000 | South, Central, North | Old World Tropics | |
| Morinda royoc | | | | | | | |
| Mucuna pruriens | No | Mesic | <1 | 101-1,000 | South | Asia | |
| Muhlenbergia capillaris | | | | | | | |
| Musa x paradisiaca | No | Mesic | 101-1,000 | 1,001-10,000 | South, Central | Asia | |
| Mynica cerifera | | | | | | | |
| Najas guadalupensis | | | | | | | |
| Nama jamaicense | | | | | | | |
| Nephrolepis multiflora | Yes | Mesic | <1 | 1,001-10,000 | South, Central | India | B 1227 |
| Neyraudia reynaudiana | Yes | Mesic | 1-100 | 1,001-10,000 | South | Southern Asia | B 1195 |
| Nymphaeoides aquatica | | | | | | | |

Frog Pond - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|--------------|--------------|---------------|------------------|---------|
| <i>Ocimum campechianum</i> | | | | | | | |
| <i>Ocolea conifera</i> | | | | | | | |
| <i>Odontosoria clavata</i> | | | | | | | |
| <i>Oecocladia maculata</i> | Yes | Mesic | <1 | 10†-1,000 | South | Old World | |
| <i>Osmunda regalis</i> var. <i>speciosissima</i> | | | | | | | |
| <i>Oxalis corniculata</i> | | | | | | | |
| <i>Panicum dichotomiflorum</i> var. <i>bartowense</i> | | | | | | | |
| <i>Panicum dichotomiflorum</i> var. <i>dichotomiflorum</i> | | | | | | | |
| <i>Panicum hemitomon</i> | | | | | | | |
| <i>Panicum maximum</i> | No | Mesic | 1-100 | 1,001-10,000 | Throughout | Africa | |
| <i>Panicum repens</i> | Yes | Wet | 10†-1,000 | >100,000 | Throughout | Tropical America | |
| <i>Panicum rigidulum</i> | | | | | | | |
| <i>Panicum virgatum</i> | | | | | | | |
| <i>Parthenium hysterophorus</i> | No | Mesic | 1,001-10,000 | >100,000 | Throughout | Tropical America | BW 424 |
| <i>Pathenocissus quinquefolia</i> | | | | | | | |
| <i>Paspalidium geminatum</i> | | | | | | | |
| <i>Paspalum blodgettii</i> | | | | | | | |
| <i>Paspalum caespitosum</i> | | | | | | | |
| <i>Paspalum conjugatum</i> | | | | | | | |
| <i>Paspalum distichum</i> | | | | | | | |
| <i>Paspalum fimbriatum</i> | No | Mesic | <1 | 1,001-10,000 | South | Tropical America | BW 357 |
| <i>Paspalum monostachyum</i> | | | | | | | |
| <i>Paspalum urvillei</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | South America | |
| <i>Passiflora suberosa</i> | | | | | | | |
| <i>Factis glaucescens</i> | | | | | | | |
| <i>Pennisetum purpureum</i> | No | Mesic | 10†-1,000 | >100,000 | Throughout | Africa | |
| <i>Pentalinon luteum</i> | | | | | | | |
| <i>Persea palustris</i> | | | | | | | |
| <i>Phlebodium aureum</i> | | | | | | | |
| <i>Phragmites australis</i> | | | | | | | |
| <i>Phyllo nodiflora</i> | | | | | | | |
| <i>Phyllo stoechadifolia</i> | | | | | | | |
| <i>Phyllanthus amarus</i> | | | | | | | |
| <i>Phyllanthus carolinensis</i> ssp. <i>saxicola</i> | | | | | | | |
| <i>Phyllanthus pentaphyllus</i> var. <i>floridanus</i> | | | | | | | |
| <i>Phyllanthus urinaria</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Old World | BW 365 |
| <i>Physalis angulata</i> | | | | | | | |
| <i>Physalis walteri</i> | | | | | | | |
| <i>Physotacca americana</i> | | | | | | | |
| <i>Pilea microphylla</i> | | | | | | | |
| <i>Pinguicula pumila</i> | | | | | | | |
| <i>Pinus elliottii</i> var. <i>densa</i> | | | | | | | |
| <i>Pithecellobium caroliniana</i> | | | | | | | |
| <i>Plantago major</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Eurasia | |
| <i>Pithecellobium carolinianum</i> | | | | | | | |
| <i>Pithecellobium floridanum</i> | | | | | | | |
| <i>Pithecellobium tenuifolium</i> | | | | | | | |
| <i>Pithecellobium torreyi</i> | | | | | | | |
| <i>Pithecellobium urbanum</i> | | | | | | | |
| <i>Pithecellobium umbellatum</i> | | | | | | | |
| <i>Pithecellobium usneoides</i> | | | | | | | |
| <i>Pithecellobium woodii</i> | | | | | | | |
| <i>Pithecellobium williamsii</i> | | | | | | | |
| <i>Pithecellobium warrenii</i> | | | | | | | |
| <i>Pithecellobium yucatanicum</i> | | | | | | | |
| <i>Pithecellobium zygomaticum</i> | | | | | | | |
| <i>Pithecellobium zygophyllum</i> | | | | | | | |
| <i>Pithecellobium</i> sp. | | | | | | | |

Frog Pond - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|-----------|----------------|---------------------------|-------------------------|---------|
| <i>Plucea rosea</i> | | | | | | | |
| <i>Poinsettia cyathophylla</i> | | | | | | | |
| <i>Poinsettia heterophylla</i> | | | | | | | |
| <i>Polygonia gracilliflora</i> | | | | | | | |
| <i>Polygonum hydropiperoides</i> | | | | | | | |
| <i>Polygonum lapathifolium</i> | No | Mesic | 101-1,000 | >100,000 | South, Central, Panhandle | Tropical America | |
| <i>Polygonum pensylvanicum</i> | No | Mesic | <1 | 1-100 | South, Central, North | Temperate North America | B 1221 |
| <i>Polygonum perfoliata</i> | No | Mesic | 101-1,000 | >100,000 | South, North | Europe | BW 421 |
| <i>Polygonum punctatum</i> | | | | | | | |
| <i>Polypondium polypodioides</i> var. <i>mitchauxiana</i> | | | | | | | |
| <i>Polypteron procumbens</i> | | | | | | | |
| <i>Pontederia cordata</i> | | | | | | | |
| <i>Portulaca oleracea</i> | No | Mesic | <1 | 10,001-100,000 | Throughout | Old World | |
| <i>Proserpinaca palustris</i> | | | | | | | |
| <i>Proserpinaca pectinata</i> | | | | | | | |
| <i>Psidium guajava</i> | Yes | Mesic | <1 | 1,001-10,000 | South, Central | Tropical America | |
| <i>Psilothrix nudum</i> | | | | | | | |
| <i>Psychotria nervosa</i> | | | | | | | |
| <i>Pteridium aquilinum</i> var. <i>caudatum</i> | | | | | | | |
| <i>Pteris bahamensis</i> | | | | | | | |
| <i>Pteris vittata</i> | Yes | Mesic | 101-1,000 | 10,001-100,000 | Throughout | Eastern Asia | BW 347 |
| <i>Ptilimnium capillaceum</i> | | | | | | | |
| <i>Quercus virginiana</i> | | | | | | | |
| <i>Randia aculeata</i> | | | | | | | |
| <i>Rapanea punctata</i> | | | | | | | |
| <i>Rhus copallina</i> | | | | | | | |
| <i>Rhynchoslytrum repens</i> | No | Mesic | 101-1,000 | >100,000 | Throughout | South Africa | B 1177 |
| <i>Rhynchosia minima</i> | | | | | | | |
| <i>Rhynchospora colorata</i> | | | | | | | |
| <i>Rhynchospora divergens</i> | | | | | | | |
| <i>Rhynchospora floridensis</i> | | | | | | | |
| <i>Rhynchospora microcarpa</i> | | | | | | | |
| <i>Rhynchospora odorata</i> | | | | | | | |
| <i>Ricinus communis</i> | Yes | Mesic | 1-100 | 101-1,000 | Throughout | Africa | |
| <i>Rivina humilis</i> | | | | | | | |
| <i>Rorippa palustris</i> spp. <i>femoriflora</i> | | | | | | | |
| <i>Rothboellia cochinchinensis</i> | No | Mesic | 1-100 | 10,001-100,000 | Throughout | Tropical Asia | B 1185 |
| <i>Ruellia succulenta</i> | | | | | | | |
| <i>Sabal palmetto</i> | | | | | | | |
| <i>Saccharum giganteum</i> | | | | | | | |
| <i>Sagittaria lancifolia</i> | | | | | | | |
| <i>Salix caroliniana</i> | | | | | | | |
| <i>Salvia serotina</i> | | | | | | | |
| <i>Sambucus canadensis</i> | | | | | | | |
| <i>Samolus ebracteatus</i> | | | | | | | |
| <i>Sarcostemma clausum</i> | | | | | | | |

Frog Pond - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range |
|--|------------|---------|-----------|----------------|---------------------------|--|
| | | Mesic | 101-1,000 | >100,000 | South, Central, North | Brazil and Paraguay |
| <i>Schinus terebinthifolius</i> | Yes | | | | | |
| <i>Schizachyrium gracile</i> | | | | | | |
| <i>Schizachyrium rhizomatum</i> | | | | | | |
| <i>Schizachyrium sanguineum</i> | | | | | | |
| <i>Schoenius nigricans</i> | | | | | | |
| <i>Schoepfia chrysophylloides</i> | | | | | | |
| <i>Selaginella eatonii</i> | | | | | | |
| <i>Senna ligustrina</i> | | | | | | |
| <i>Senna obtusifolia</i> | | | | | | |
| <i>Serenoa repens</i> | | | | | | |
| <i>Sesbania herbacea</i> | | | | | | |
| <i>Setaria conuta</i> | | | | | | |
| <i>Setaria parviflora</i> | | | | | | |
| <i>Sida acuta</i> | | | | | | |
| <i>Sida rhombifolia</i> | | | | | | |
| <i>Sideroxylon declinatum</i> ssp. <i>austrofloridense</i> | | | | | | |
| <i>Sideroxylon salicifolium</i> | | | | | | |
| <i>Simarouba glauca</i> | | | | | | |
| <i>Smilax auriculata</i> | | | | | | |
| <i>Smilax bona-nox</i> | | | | | | |
| <i>Smilax laurifolia</i> | | | | | | |
| <i>Solanum americanum</i> | | | | | | |
| <i>Solanum verbascoifolium</i> | | | | | | |
| <i>Solidago gigantea</i> | | | | | | |
| <i>Solidago odora</i> var. <i>chapmanii</i> | | | | | | |
| <i>Solidago sempervirens</i> | | | | | | |
| <i>Solidago stricta</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Europe |
| <i>Sonchus oleraceus</i> | No | Mesic | <1 | 1-100 | Throughout | Europe |
| <i>Sorghum arundinaceum</i> | No | Mesic | 101-1,000 | 10,001-100,000 | South, Central, Panhandle | Europe |
| <i>Sorghum bicolor</i> | No | Mesic | 101-1,000 | 10,001-100,000 | Throughout | Old World |
| <i>Spartina bakeri</i> | | | | | | |
| <i>Spermatoceae assurgens</i> | | | | | | |
| <i>Spermatoceae floridana</i> | | | | | | |
| <i>Spermatoceae prostrata</i> | | | | | | |
| <i>Spermatoceae terminalis</i> | | | | | | |
| <i>Spermatoceae tetraquetria</i> | Yes | Mesic | 101-1,000 | >100,000 | South, Central, North | Africa |
| <i>Spermatoceae verticillata</i> | | | | | | |
| <i>Spigelia anthelmia</i> | | | | | | |
| <i>Sporobolus indicus</i> var. <i>pyramidalis</i> | No | Mesic | 101-1,000 | >100,000 | South, Central, Panhandle | Tropical America |
| <i>Stachytaphila jamaicensis</i> | | | | | | |
| <i>Stenandrium duce</i> | | | | | | |
| <i>Swietenia mahagoni</i> | No | Mesic | <1 | 1-100 | South | Extreme southern Florida, West Indies, Central and South America |
| <i>Tabea bauera</i> | | | | | | |
| <i>Tetrazygia bicolor</i> | | | | | | |
| <i>Teucrium canadense</i> | | | | | | |

Frog Pond - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|-----------|--------------|---------------------------|------------------------------|---------|
| <i>Thalia geniculata</i> | | | | | | | |
| <i>Thelypteris kunthii</i> | | | | | | | |
| <i>Thelypteris ovala</i> | | | | | | | |
| <i>Thelypteris palustris</i> var. <i>pubescens</i> | | | | | | | |
| <i>Tillandsia balbisiana</i> | | | | | | | |
| <i>Tillandsia fasciculata</i> var. <i>densispica</i> | | | | | | | |
| <i>Tillandsia setacea</i> | | | | | | | |
| <i>Tillandsia utricularia</i> | | | | | | | |
| <i>Toxicodendron radicans</i> | | | | | | | |
| <i>Trema lamarckianum</i> | | | | | | | |
| <i>Trema micranthum</i> | | | | | | | |
| <i>Trichostema dichotomum</i> | | | | | | | |
| <i>Tridax procumbens</i> | Yes | Mesic | 101-1,000 | 1,001-10,000 | South, Central | Tropical America | BW 348 |
| <i>Typha domingensis</i> | | | | | | | |
| <i>Urena lobata</i> | | | | | | | |
| <i>Urochloa adspersa</i> | | | | | | | |
| <i>Urochloa mutica</i> | Yes | Wet | 1-100 | 1,001-10,000 | South, Central | | B 1211 |
| <i>Urochloa plantaginea</i> | No | Mesic | <1 | 101-1,000 | South, Central, Panhandle | Mexico to Bolivia and Brazil | BW 427 |
| <i>Urucumaria foliosa</i> | | | | | | | |
| <i>Urucumaria purpurea</i> | | | | | | | |
| <i>Verbena bonariensis</i> | No | Mesic | <1 | 101-1,000 | Throughout | South America | |
| <i>Verbena brasiliensis</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | South America | |
| <i>Verbena scabra</i> | | | | | | | |
| <i>Vigna luteola</i> | | | | | | | |
| <i>Vitis cinerea</i> var. <i>floridana</i> | | | | | | | |
| <i>Vitis rotundifolia</i> | | | | | | | |
| <i>Waltheria indica</i> | | | | | | | |
| <i>Washingtonia robusta</i> | No | Mesic | <1 | 1-100 | South, Central | Mexico | |
| <i>Zamia integrifolia</i> | | | | | | | |

Frog Pond - Endangered and Threatened Taxa

| | State | Habitat | Acreage | Pop. Size | Status | Stations | Natural Range | Voucher |
|--|-------|---------|----------|--------------|-------------|------------|---|---------|
| <i>Anemia wrightii</i> | E | Mesic | <1 | 101-1,000 | Stable | | 2 Florida, Cuba | BW 1204 |
| <i>Bletia purpurea</i> | T | Mesic | <1 | 1-100 | Decreasing? | | 1 Florida, West Indies, Central and South America | |
| <i>Byrsinima lucida</i> | E | Mesic | 1-100 | 101-1,000 | Stable | | 2 Florida and West Indies | |
| <i>Eulophia alta</i> | T | Wet | <1 | 1-100 | Increasing? | | 1 Florida, West Indies, Central and South America | |
| <i>Ilex krugiana</i> | E | Mesic | 1-100 | 1-100 | Increasing? | | 1 Florida, Bahamas, Cuba | |
| <i>Jacquemontia curtissii</i> | E | Mesic | 1-100 | 101-1,000 | Stable | | 2 South Florida | BW 423 |
| <i>Melanthera parvifolia</i> | E | Mesic | 101-1,00 | 1,001-10,000 | Stable | Throughout | South Florida | |
| <i>Odontosoria clavata</i> | E | Mesic | 1-100 | 1,001-10,000 | Stable | | 4 Florida, Bahamas, Greater Antilles, S. Mexico | B 1198 |
| <i>Selaginella eatonii</i> | E | Mesic | 1-100 | 1,001-10,000 | Stable | Throughout | Florida, Bahamas | B 1205 |
| <i>Tetrazygia bicolor</i> | T | Mesic | 1-100 | 101-1,000 | Stable | | 2 Florida, Bahamas, Cuba | |
| <i>Tillandsia balbisiana</i> | T | Mesic | 1-100 | 101-1,000 | Stable | Throughout | Florida to South America | |
| <i>Tillandsia fasciculata</i> var. <i>densispica</i> | E | Mesic | 1-100 | 101-1,000 | Stable | Throughout | Florida to Central America | |
| <i>Tillandsia utriculata</i> | E | Mesic | 1-100 | 101-1,000 | Stable | Throughout | Georgia to South America | |

Lake Okeechobee Marshes - Preliminary List of Vascular Plants

Bold = Exotic; Italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|----------------|----------------|---------------------------|------------------|----------------|
| <i>Abrus precatorius</i> | No | Mesic | <1 | 101-1,000 | South, Central | Old World | |
| <i>Acer rubrum</i> | | | | | | | |
| <i>Acrostichum danaeifolium</i> | | | | | | | |
| <i>Alternanthera philoxeroides</i> | Yes | Wet | 10,001-100,000 | >100,000 | Throughout | South America | |
| <i>Alternanthera paronychoides</i> | No | Wet | <1 | 1-100 | South, Central, Panhandle | Tropical America | BW 405A |
| <i>Amaranthus australis</i> | | | | | | | |
| <i>Amaranthus viridis</i> | No | Mesic | <1 | 1-100 | Throughout | Tropical America | |
| <i>Ambrosia artemisiifolia</i> | | | | | | | |
| <i>Ammannia latifolia</i> | | | | | | | |
| <i>Ampelopsis arborea</i> | | | | | | | |
| <i>Annona glabra</i> | | | | | | | |
| <i>Baccharis halimifolia</i> | | | | | | | |
| <i>Bacopa monnieri</i> | | | | | | | |
| <i>Bambusa vulgaris</i> | No | Mesic | <1 | 1-100 | South, Central | Old World | |
| <i>Bidens laevis</i> | No | Mesic | 1-100 | 101-1,000 | South, Central | Tropical Asia | GB 989 |
| <i>Bischofia javanica</i> | | | | | | | |
| <i>Boehmeria cylindrica</i> | | | | | | | |
| <i>Canna flaccida</i> | | | | | | | |
| <i>Carica papaya</i> a | Yes | Mesic | 1-100 | 1,001-10,000 | South, Central | Tropical America | GB 969 |
| <i>Casuarina equisetifolia</i> | No | Mesic | 101-1,000 | 10,001-100,000 | South, Central, North | Australia | |
| <i>Casuarina glauca</i> | No | Mesic | 101-1,000 | 10,001-100,000 | South, Central | Australia | |
| <i>Catharanthus roseus</i> | No | Mesic | 1-100 | 1-100 | South, Central, Panhandle | Madagascar | GB 967 |
| <i>Cattil laevigata</i> | | | | | | | |
| <i>Centella asiatica</i> | | | | | | | |
| <i>Cephaelanthus occidentalis</i> | | | | | | | |
| <i>Ceratophyllum demersum</i> | No | Mesic | 1-100 | 101-1,000 | South, Central | West Indies | GB 955 |
| <i>Cestrum diurnum</i> | | | | | | | |
| <i>Chamaesyce cordifolia</i> | | | | | | | |
| <i>Cicuta maculata</i> | | | | | | | |
| <i>Cissus verticillata</i> | | | | | | | |
| <i>Citrus x paradisi</i> | No | Mesic | 1-100 | 101-1,000 | South, Central | Southeast Asia | GB 1015 |
| <i>Cladium jamaicense</i> | | | | | | | |
| <i>Coccoloba uvifera</i> | | | | | | | |
| <i>Colocasia esculenta</i> | Yes | Wet | 1,001-10,000 | 10,001-100,000 | Throughout | Asia | |
| <i>Commelinia diffusa</i> | | | | | | | |
| <i>Commelinia diffusa</i> var. <i>gigas</i> | | | | | | | |
| <i>Conyza canadensis</i> var. <i>pusilla</i> | | | | | | | |
| <i>Crinum</i> sp. | No | Mesic | <1 | 1-100 | Unknown | Unknown | |
| <i>Crinum glandulosus</i> | | | | | | | |
| <i>Cucurbita okeechobeensis</i> | | | | | | | |

Lake Okeechobee Marshes - Preliminary List of Vascular Plants

Bold = Exotic; *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|----------------------------------|------------|---------|----------------|-----------|----------------|---|---------|
| Cupaniopsis anacardioides | No | Mesic | 1-100 | 100-1,000 | South, Central | Australia | GB 980 |
| <i>Cynanchum scoparium</i> | No | Mesic | 1-100 | 100-1,000 | Throughout | Africa | |
| Cynodon dactylon | No | Mesic | 1-100 | 100-1,000 | Throughout | | |
| <i>Cyperus compressus</i> | | | | | | | |
| <i>Cyperus croceus</i> | | | | | | | |
| <i>Cyperus diffusus</i> | | | | | | | |
| <i>Cyperus flavescens</i> | | | | | | | |
| <i>Cyperus odoratus</i> | | | | | | | |
| <i>Cyperus polystachyos</i> | | | | | | | |
| <i>Cyperus surinamensis</i> | | | | | | | |
| <i>Dicliptera sexangularis</i> | | | | | | | |
| <i>Digitaria ciliaris</i> | | | | | | | |
| <i>Echinochloa walteri</i> | | | | | | | |
| <i>Ecliptia prostrata</i> | | | | | | | |
| Eichhornia crassipes | Yes | Wet | 10,001-100,000 | >100,000 | Throughout | Brazil | |
| <i>Eleocharis cellulosa</i> | | | | | | | |
| <i>Eleocharis geniculata</i> | | | | | | | |
| <i>Eleocharis interstincta</i> | | | | | | | |
| Eleusine indica | No | Mesic | <1 | 1-100 | Throughout | Old World | GB 996 |
| <i>Erechtites hieracifolia</i> | | | | | | | |
| <i>Eugenia uniflora</i> | Yes | Mesic | <1 | 1-100 | South, Central | South America | GB 1001 |
| <i>Eupatorium capillifolium</i> | | | | | | | |
| <i>Eupatorium serotinum</i> | | | | | | | |
| <i>Eustachys petraea</i> | | | | | | | |
| Ficus altissima | No | Mesic | <1 | 1-100 | South | Assam to Malay Archipelago, Philippines | |
| <i>Ficus aurea</i> | | | | | | | |
| Ficus microcarpa | No | Mesic | 1-100 | 101-1,000 | South, Central | Old World | GB 984 |
| <i>Fimbristylis autumnalis</i> | | | | | | | |
| <i>Forestiera segregata</i> | | | | | | | |
| <i>Fraxinus caroliniana</i> | | | | | | | |
| <i>Habenaria repens</i> | | | | | | | |
| <i>Helianthus agrestis</i> | | | | | | | |
| <i>Hibiscus grandiflorus</i> | | | | | | | |
| Hydroilla verticillata | Yes | Wet | 10,001-100,000 | >100,000 | Throughout | Old World | BW 408 |
| <i>Hydrocotyle umbellata</i> | | | | | | | |
| <i>Hydrocotyle verticillata</i> | | | | | | | |
| <i>Ipomoea alba</i> | | | | | | | |
| <i>Iresine diffusa</i> | | | | | | | |
| <i>Kosteletzya virginica</i> | | | | | | | |
| <i>Kyllinga brevifolia</i> | | | | | | | |
| <i>Lactuca floridana</i> | | | | | | | |
| <i>Lemna valdiviana</i> | | | | | | | |
| <i>Limnobium spongia</i> | | | | | | | |

Lake Okeechobee Marshes - Preliminary List of Vascular Plants

Bold = Exotic; *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|----------------|-----------|-----------------------|-------------------|---------|
| Ludwigia octovalvis | | | | | | | |
| <i>Ludwigia peploides</i> | | | | | | | |
| <i>Ludwigia repens</i> | | | | | | | |
| <i>Lytium alatum</i> var. <i>lanceolatum</i> | | | | | | | |
| Macrothelypteris torresiana | No | Mesic | <1 | 1-100 | Throughout | Asia | BW 397 |
| Malealeuca quinquenervia | Yes | Wet | 10,001-100,000 | >100,000 | South, Central | Australia | GB 994 |
| <i>Melothria pendula</i> | | | | | | | |
| <i>Menzelia floridana</i> | | | | | | | |
| <i>Mikania scandens</i> | | | | | | | |
| Mimordica charantia | No | Mesic | 1-100 | 101-1,000 | South, Central, North | Old World Tropics | GB 1003 |
| <i>Morus rubra</i> | | | | | | | |
| Musa x paradisiaca | No | Mesic | <1 | 1-100 | South, Central, North | Asia | |
| <i>Myrica cerifera</i> | | | | | | | |
| <i>Najas guadalupensis</i> | | | | | | | |
| <i>Najas cf. minor</i> | Yes | Wet | 1-100 | 100-1,000 | Panhandle | ? | BW 391 |
| <i>Neelumbo lutea</i> | | | | | | | |
| <i>Nuphar lutea</i> ssp. <i>advena</i> | | | | | | | |
| <i>Nymphaea mexicana</i> | | | | | | | |
| <i>Nymphaea odorata</i> | | | | | | | |
| <i>Nymphaea x thiona</i> | | | | | | | |
| <i>Ophiolismenus heterotilus</i> | | | | | | | |
| <i>Panicum dichotomiflorum</i> var. <i>bartowense</i> | | | | | | | |
| <i>Panicum hemitonon</i> | | | | | | | |
| Panicum repens | Yes | Wet | 1,000-10,000 | >100,000 | Throughout | Tropical America | |
| <i>Paspalidium geminatum</i> | | | | | | | |
| <i>Paspalum conjugatum</i> | | | | | | | |
| <i>Paspalum repens</i> | | | | | | | |
| <i>Passiflora suberosa</i> | | | | | | | |
| <i>Peltandra virginica</i> | | | | | | | |
| Pennisetum purpureum | No | Wet | 1,000-10,000 | >100,000 | Throughout | Africa | |
| Persea americana | No | Mesic | <1 | 1-100 | South, Central | Tropical America | GB 104 |
| <i>Peltiveria alliacea</i> | | | | | | | |
| <i>Phlebodium aureum</i> | | | | | | | |
| <i>Phragmites australis</i> | | | | | | | |
| <i>Phyla nodiflora</i> | | | | | | | |
| <i>Phytolacca americana</i> | | | | | | | |
| <i>Pisonia aculeata</i> | | | | | | | |
| Pistia stratiotes | Yes | Wet | 10,001-100,000 | >100,000 | Throughout | Tropical America | |
| <i>Pluchea odorata</i> | | | | | | | |
| <i>Poinsettia cyathophora</i> | | | | | | | |
| <i>Polygonum densiflorum</i> | | | | | | | |
| <i>Polygonum hydropiperoides</i> | | | | | | | |
| <i>Polygonum punctatum</i> | | | | | | | |

Lake Okeechobee Marshes - Preliminary List of Vascular Plants

Bold = Exotic; *Italics* = Cultivated

| Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|---------|---------|-----------|----------------|-----------------------|------------------------------------|
| <i>Polypterygium procumbens</i> | | | | | | |
| <i>Pontederia cordata</i> | | | | | | |
| <i>Potamogeton illinoensis</i> | | | | | | |
| <i>Potamogeton pusillus</i> | | | | | | |
| <i>Psidium guajava</i> | Yes | Mesic | 1-100 | 1,001-10,000 | South, Central | Tropical America |
| <i>Pieris tripartita</i> | No | Mesic | <1 | 1-100 | South, Central | Tropical Asia |
| <i>Pieris virgata</i> | No | Mesic | <1 | 1-100 | Throughout | Eastern Asia |
| <i>Quercus virginiana</i> | | | | | | |
| <i>Rhynchospora nitens</i> | | | | | | |
| <i>Ricinus communis</i> | Yes | Mesic | 101-1,000 | 1,001-10,000 | Throughout | Africa |
| <i>Rivina humilis</i> | No | Mesic | <1 | 1-100 | South | Florida, Bahamas, Cuba, Hispaniola |
| <i>Roxystonea regia</i> | | | | | | |
| <i>Rumex verticillatus</i> | | | | | | |
| <i>Sabal palmetto</i> | | | | | | |
| <i>Sagittaria lancifolia</i> | | | | | | |
| <i>Sagittaria latifolia</i> | | | | | | |
| <i>Sagittaria cf. filiformis</i> | | | | | | |
| <i>Salix caroliniana</i> | | | | | | |
| <i>Salvia occidentalis</i> | | | | | | |
| <i>Salvinia minima</i> | | | | | | |
| <i>Sambucus canadensis</i> | | | | | | |
| <i>Sarcostemma clausum</i> | | | | | | |
| <i>Saururus cernuus</i> | | | | | | |
| <i>Schinus terebinthifolius</i> | Yes | Mesic | 101-1,000 | 10,001-100,000 | South, Central, North | Brazil and Paraguay |
| <i>Scirpus pungens</i> | | | | | | |
| <i>Scirpus tabernaemontani</i> | | | | | | |
| <i>Scoparia dulcis</i> | | | | | | |
| <i>Senna alata</i> | No | Wet | <1 | 1-100 | South, Central | Tropical America |
| <i>Senna obtusifolia</i> | | | | | | |
| <i>Senna pendula var. glabrata</i> | No | Mesic | <1 | 1-100 | South, Central | South America |
| <i>Sesbania herbacea</i> | | | | | | |
| <i>Setaria magna</i> | | | | | | |
| <i>Setaria parviflora</i> | | | | | | |
| <i>Sida acuta</i> | | | | | | |
| <i>Sida hombifolia</i> | | | | | | |
| <i>Smilax tamnoides</i> | | | | | | |
| <i>Solanum capsicoides</i> | | | | | | |
| <i>Solanum cf. torvum</i> | Yes | Mesic | <1 | 1-100 | South, Central, North | West Indies |
| <i>Sonchus oleraceus</i> | No | Mesic | <1 | 1-100 | Throughout | Europe |
| <i>Spartina bakeri</i> | | | | | | |
| <i>Spirodela polyrhiza</i> | | | | | | |
| <i>Sporopholus indicus</i> var. <i>indicus</i> | No | Mesic | <1 | 1-100 | Throughout | Tropical Asia |
| <i>Syzygium cumini</i> | No | Mesic | 1-100 | 1-100 | South | Asia |

Lake Okeechobee Marshes - Preliminary List of Vascular Plants

Bold = Exotic; /italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|-------------------------------|------------|---------|---------|-----------|----------------|---------------|---------|
| <i>Taxodium ascendens</i> | | | | | | | |
| <i>Taxodium distichum</i> | | | | | | | |
| <i>Tectaria incisa</i> | | | | | | | |
| <i>Thalia geniculata</i> | | | | | | | |
| <i>Thelypteris kunthii</i> | | | | | | | |
| <i>Tillandsia recurvata</i> | | | | | | | |
| <i>Tillandsia usneoides</i> | | | | | | | |
| <i>Tillandsia utriculata</i> | | | | | | | |
| <i>Toxicodendron radicans</i> | | | | | | | |
| <i>Typha domingensis</i> | No | Mesic | <1 | 1-100 | South, Central | Asia | GB 997 |
| <i>Urochloa subquadripara</i> | | | | | | | |
| <i>Utricularia foliosa</i> | | | | | | | |
| <i>Utricularia purpurea</i> | | | | | | | |
| <i>Vallisneria americana</i> | | | | | | | |
| <i>Verbena scabra</i> | | | | | | | |
| <i>Vigna luteola</i> | | | | | | | |
| <i>Washingtonia robusta</i> | No | Mesic | <1 | 1-100 | South, Central | Mexico | GB 982 |
| <i>Youngia japonica</i> | No | Mesic | <1 | 1-100 | Throughout | SE Asia | |
| <i>Zizaniopsis miliacea</i> | | | | | | | |

Lake Okeechobee Marshes- Endangered and Threatened Taxa

| | State Status | Habitat | Pop. Size | # Stations | Status | Natural Range | Voucher |
|---------------------------------|--------------|---------|-----------|------------|--------|---------------|--|
| <i>Cucurbita okeechobeensis</i> | E | Wet | 1-100 | 101-1,000 | 7 | Increasing? | Central Florida |
| <i>Tillandsia utriculata</i> | E | Mesic | <1 | 1-100 | 1 | Stable | Georgia and Florida, West Indies, Mexico, Northern South America, Venezuela BW 383, 389, 393, 398 |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|---|------------|---------|---------|--------------|----------------|-----------------------------------|---------|
| <i>Abrus precatorius</i> | Yes | Mesic | <1 | 1-100 | South, Central | Old World | BW 734 |
| <i>Acacia auriculiformis</i> | No | Mesic | <1 | 1-100 | South | Australia | BW 728 |
| <i>Acalypha gracilens</i> | | | | | | | |
| <i>Acer rubrum</i> | | | | | | | |
| <i>Acrostichum danaeifolium</i> | | | | | | | |
| <i>Aeschynomene americana</i> | | | | | | | |
| <i>Agalinis fasciculata</i> | | | | | | | |
| <i>Agalinis linifolia</i> | | | | | | | |
| <i>Ageratina jucunda</i> | | | | | | | |
| <i>Ageratum houstonianum</i> | No | Mesic | <1 | 1-100 | South, Central | Mexico and northern South America | BW 730 |
| <i>Albizia lebbeck</i> | No | Mesic | <1 | 1-100 | South, Central | Asia | |
| <i>Alnus lutea</i> | | | | | | | |
| <i>Alamanda cathartica</i> | No | Mesic | <1 | 1-100 | South, Central | Tropical America | BW 60 |
| <i>Alternanthera philoxeroides</i> | Yes | Wet | 1-100 | 1,001-10,000 | Throughout | South America | |
| <i>Alternanthera sessilis</i> | Yes | Wet | <1 | 1-100 | Throughout | Pantropical | |
| <i>Amaranthus spinosus</i> | No | Mesic | <1 | 101-1,000 | Throughout | India | BW 586 |
| <i>Amaranthus viridis</i> | No | Mesic | <1 | 101-1,000 | Throughout | Tropical America | BW 630 |
| <i>Ambrosia artemisiifolia</i> | | | | | | | |
| <i>Anmannia coccinea</i> | | | | | | | |
| <i>Annona latifolia</i> | | | | | | | |
| <i>Anopha fruticosa</i> | | | | | | | |
| <i>Ampeleopsis atropurpurea</i> | | | | | | | |
| <i>Amphicarpum muhlenbergianum</i> | | | | | | | |
| <i>Andropogon brachystachyus</i> | | | | | | | |
| <i>Andropogon floridanus</i> | | | | | | | |
| <i>Andropogon glomeratus</i> var. <i>glaucoptis</i> | | | | | | | |
| <i>Andropogon glomeratus</i> var. <i>hirsutior</i> | | | | | | | |
| <i>Andropogon glomeratus</i> var. <i>pumilus</i> | | | | | | | |
| <i>Andropogon gyrans</i> | | | | | | | |
| <i>Andropogon tenuarius</i> | | | | | | | |
| <i>Andropogon virginicus</i> var. <i>decipiens</i> | | | | | | | |
| <i>Andropogon virginicus</i> var. <i>glaucus</i> | | | | | | | |
| <i>Andropogon virginicus</i> var. <i>virginicus</i> | | | | | | | |
| <i>Annona glabra</i> | | | | | | | |
| <i>Aplospadix americana</i> | | | | | | | |
| <i>Aperta aphylla</i> | | | | | | | |
| <i>Ardisia escallonioides</i> | | | | | | | |
| <i>Arisaema triphyllum</i> | | | | | | | |
| <i>Asplenium beyrichiana</i> | | | | | | | |
| <i>Ansiedia palustris</i> | | | | | | | |
| <i>Ansiedia purpurascens</i> var. <i>tenuispica</i> | | | | | | | |
| <i>Ansiedia rhizomophora</i> | | | | | | | |
| <i>Ansiedia spiciformis</i> | | | | | | | |
| <i>Asclepias lanceolata</i> | | | | | | | |
| <i>Asclepias pedicellata</i> | | | | | | | |
| <i>Asimina reticulata</i> | | | | | | | |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|---|------------|---------|---------|--------------|-----------------------|--|---------------|
| <i>Aster bracelet</i> | | | | | | | |
| <i>Aster dumosus</i> | | | | | | | |
| <i>Aster iontiifolius</i> | | | | | | | |
| <i>Axonopus fissifolius</i> | | | | | | | |
| <i>Axonopus furcatus</i> | | | | | | | |
| <i>Azolla caroliniana</i> | | | | | | | |
| <i>Baccharis glomeruliflora</i> | | | | | | | |
| <i>Baccharis halimifolia</i> | | | | | | | |
| <i>Bacopa caroliniana</i> | | | | | | | |
| <i>Bacopa monnieri</i> | | | | | | | |
| <i>Baldwinia angustifolia</i> | | | | | | | |
| <i>Bambusa vulgaris</i> | No | Mesic | <1 | 1-100 | Throughout | Asia | |
| <i>Bemania racemosa</i> | | | | | | | |
| <i>Bidens alba</i> var. <i>radiata</i> | | | | | | | |
| <i>Bigelowia nudata</i> ssp. <i>australis</i> | | | | | | | |
| <i>Bischofia javanica</i> | Yes | Mesic | <1 | 1-100 | South, Central | Tropical Asia | <i>BW 537</i> |
| <i>Blechnum serratum</i> | | | | | | | |
| <i>Blechnum pyramidatum</i> | No | Mesic | <1 | 1,001-10,000 | South, Central, North | Tropical America | |
| <i>Boehmeria cylindrica</i> | | | | | | | |
| <i>Bulbostylis barbata</i> | No | Mesic | <1 | 1-100 | Throughout | Old World Tropics | <i>BW 69</i> |
| <i>Bulbostylis ciliatifolia</i> | | | | | | | |
| <i>Bulbostylis stenophylla</i> | | | | | | | |
| <i>Burmannia capitata</i> | | | | | | | |
| <i>Callicarpa americana</i> | | | | | | | |
| <i>Callisia fragrans</i> | No | Mesic | <1 | 1-100 | South, Central | Mexico | <i>BW 63</i> |
| <i>Carex stipigera</i> ssp. <i>limnophila</i> | | | | | | | |
| <i>Campyloneuron phyllidis</i> | | | | | | | |
| <i>Canna flaccida</i> | | | | | | | |
| <i>Cardamine pensylvanica</i> | | | | | | | |
| <i>Carex lupulina</i> | | | | | | | |
| <i>Carex longii</i> | | | | | | | |
| <i>Carica papaya</i> | Yes | Mesic | <1 | 101-1,000 | South, Central | Tropical America | |
| <i>Caprhephorus camosus</i> | | | | | | | |
| <i>Caprhephorus corymbosus</i> | | | | | | | |
| <i>Caprhephorus odoratissimus</i> | | | | | | | |
| <i>Cayya aquatica</i> | | | | | | | |
| <i>Caryota mitis</i> | No | Mesic | <1 | 1-100 | South | Burma to Malay Pen., Java, and Philippines | |
| <i>Cassytha filiformis</i> | | | | | | | |
| <i>Casuarina equisetifolia</i> | No | Mesic | 1-100 | 101-1,000 | South, Central, North | Australia | |
| <i>Cenchrus echinatus</i> | | | | | | | |
| <i>Cenchrus incertus</i> | | | | | | | |
| <i>Centella asiatica</i> | | | | | | | |
| <i>Cephalanthus occidentalis</i> | | | | | | | |
| <i>Ceratiola ericoides</i> | | | | | | | |
| <i>Ceratophyllum demersum</i> | | | | | | | |
| <i>Chamaecrista fasciata</i> | | | | | | | |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|---|---------|---------|------------|---------------|-----------------------|----------------------------|
| <i>Chamaecrista nictitans</i> var. <i>aspera</i> | | | | | | |
| <i>Chamaesyce blodgettii</i> | | | | | | |
| <i>Chamaesyce hirta</i> | | | | | | |
| <i>Chamaesyce hypericifolia</i> | | | | | | |
| <i>Chamaesyce hyssopifolia</i> | | | | | | |
| <i>Chaptalia tomentosa</i> | | | | | | |
| <i>Chenopodium ambrosioides</i> | No | Mesic | <1 | 101-1,000 | Throughout | Tropical America |
| <i>Chromolaena odorata</i> | | | | | | |
| <i>Chrysidocarpus lutescens</i> | | | | | | |
| <i>Chrysobalanus icaco</i> | | | | | | |
| <i>Chrysophyllum oliviforme</i> | | | | | | |
| <i>Citrus horridum</i> | Yes | Mesic | 1-100 | 1,001-10,000 | Throughout | |
| <i>Citrus aurantium</i> | Yes | Mesic | 1-100 | 101-1,000 | South, Central, North | Southeast Asia |
| <i>Citrus sinensis</i> | Yes | Mesic | 1-100 | 101-1,000 | South, Central | Southeast Asia |
| <i>Citrus x paradisi</i> | Yes | Mesic | 1-100 | 101-1,000 | South, Central | Southeast Asia |
| <i>Cladium jamaicense</i> | | | | | | |
| <i>Colocasia esculenta</i> | No | Wet | 1-100 | 101-1,000 | Throughout | Asia |
| <i>Commelinia diffusa</i> | | | | | | |
| <i>Conoclinium coelestinum</i> | | | | | | |
| <i>Convya canadensis</i> var. <i>pusilla</i> | No | Mesic | <1 | 1-100 | Throughout | Asia |
| <i>Corchorus aestuans</i> | Yes | Mesic | 1-100 | 101-1,000 | South | Africa |
| <i>Cordia myxa</i> | | | | | | |
| <i>Coreopsis leavenworthii</i> | | | | | | |
| <i>Corinus foemina</i> | | | | | | |
| <i>Crinum americanum</i> | | | | | | |
| <i>Crotalaria lanceolata</i> | No | Mesic | <1 | 1-100 | Throughout | Africa |
| <i>Crotalaria pallida</i> var. <i>obovata</i> | No | Mesic | <1 | 101-1,000 | South, Central | Africa |
| <i>Crotalaria rotundifolia</i> | No | Mesic | <1 | 101-1,000 | Throughout | Old World Tropics |
| <i>Crotalaria spectabilis</i> | | | | | | |
| <i>Croton glandulosus</i> var. <i>glandulosus</i> | | | | | | |
| <i>Cupaniopsis anacardioides</i> | No | Mesic | <1 | 1-100 | South, Central | Australia |
| <i>Cuphea carthagenensis</i> | No | Wet | 1-100 | 101-1,000 | Throughout | South and Central America? |
| <i>Cuthbertia ornata</i> | | | | | | |
| <i>Cynanchum angustifolium</i> | | | | | | |
| <i>Cynanchum scoparium</i> | | | | | | |
| <i>Cynodon dactylon</i> | No | Mesic | 1-100 | 1,001-10,000 | Throughout | Africa |
| <i>Cyperus compressus</i> | | | | | | |
| <i>Cyperus croceus</i> | | | | | | |
| <i>Cyperus distictus</i> | | | | | | |
| <i>Cyperus esculentus</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Old World |
| <i>Cyperus flavescens</i> | | | | | | |
| <i>Cyperus haspan</i> | | | | | | |
| <i>Cyperus ligularis</i> | | | | | | |
| <i>Cyperus odoratus</i> | | | | | | |
| <i>Cyperus polystachyos</i> | | | | | | |
| <i>Cyperus retrorsus</i> | | | | | | |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|--------------|-----------------------|--|------------------|
| Cyperus sphacelatus | No | Mesic | 1-100 | 101-1,000 | South, Central | Old and New World Tropics | BW 618, 719, 740 |
| <i>Cyperus tetragonus</i> | No | Mesic | 1-100 | 101-1,000 | Throughout | Old World Tropics | BW 637 |
| Dactyloctenium aegyptium | No | Mesic | 1-100 | 1-100 | South | Madagascar | |
| Daea pinnata | No | Mesic | <1 | 1-100 | | | |
| Delonix regia | | | | | | | |
| <i>Desmodium incanum</i> | | | | | | | |
| <i>Desmodium paniculatum</i> | | | | | | | |
| <i>Desmodium tortuosum</i> | | | | | | | |
| Desmodium triflorum | No | Mesic | <1 | 1,001-10,000 | Throughout | Old World | |
| Dichanthelium aciculare | | | | | | | |
| <i>Dichanthelium commutatum</i> | | | | | | | |
| <i>Dichanthelium dichotomum</i> | | | | | | | |
| <i>Dichanthelium ensifolium</i> var. <i>breve</i> | | | | | | | |
| <i>Dichanthelium ensifolium</i> var. <i>ensifolium</i> | | | | | | | |
| <i>Dichanthelium erectifolium</i> | | | | | | | |
| <i>Dichanthelium laxiflorum</i> | | | | | | | |
| <i>Dichanthelium leucostachys</i> | | | | | | | |
| <i>Dichanthelium portoricense</i> | | | | | | | |
| <i>Dichanthelium strigosum</i> var. <i>glabrescens</i> | | | | | | | |
| <i>Dichondra caroliniana</i> | | | | | | | |
| <i>Digita villoso</i> | | | | | | | |
| <i>Diospyros virginiana</i> | | | | | | | |
| <i>Drosera brevifolia</i> | | | | | | | |
| <i>Drosera capillaris</i> | | | | | | | |
| <i>Drymaria cordata</i> | | | | | | | |
| <i>Dyschoriste angusta</i> | | | | | | | |
| <i>Echinocloa colonia</i> | No | Mesic | <1 | 1-100 | Throughout | Old World Tropics | BW 717 |
| <i>Echinocloa walteri</i> | | | | | | | |
| <i>Eclipta prostrata</i> | | | | | | | |
| Eichhornia crassipes | Yes | Wet | 1-100 | 1,001-10,000 | Throughout | Brazil | |
| <i>Eleocharis baldwinii</i> | | | | | | | |
| <i>Eleocharis cellulosa</i> | | | | | | | |
| <i>Eleocharis flavescens</i> | | | | | | | |
| <i>Eleocharis nigrescens</i> | Yes | Mesic | 1-100 | 1,001-10,000 | Throughout | Pantropical | BW 599 |
| <i>Elephantopus elatus</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Old World | BW 86, 540 |
| Elymus Indica | | | | | | | |
| <i>Elyonurus tripacioides</i> | | | | | | | |
| Emilia fosbergii | No | Mesic | <1 | 101-1,000 | Throughout | Old World | BW 67 |
| Emilia sonchifolia | No | Mesic | <1 | 101-1,000 | South, Central, North | Old World | BW 581 |
| <i>Encyclia tampensis</i> | | | | | | | |
| <i>Epipremnum pinnatum</i> | Yes | Wet | 1-100 | 101-1,000 | South | Philippine Islands to Malay Peninsula, and Australia | BW 62 |
| <i>Eragrostis atrovirens</i> | Yes | Mesic | 1-100 | 1,001-10,000 | Throughout | Southeast Asia | |
| <i>Eragrostis ciliaris</i> | No | Mesic | 1-100 | 1,001-10,000 | Throughout | Coastal Plain, Fl. to Tex. and N.J. - (W.I., Mex., C.A., S.A.) | BW 99 |
| <i>Eragrostis elliotii</i> | | | | | | | |
| <i>Eragrostis virginica</i> | | | | | | | |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|----------------|----------------|---------------------------------|------------|
| <i>Erichites hieracifolia</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Southeast Asia | BW 592 |
| <i>Eremochloa ophiuroides</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | | |
| <i>Erigeron quercifolius</i> | | | | | | | |
| <i>Erigeron vernis</i> | | | | | | | |
| <i>Eriocaulon decangulare</i> | | | | | | | |
| <i>Eriocaulon ravenelii</i> | | | | | | | |
| <i>Eryngium aromaticum</i> | | | | | | | |
| <i>Eryngium yuccifolium</i> | | | | | | | |
| <i>Erythrina herbacea</i> | | | | | | | |
| <i>Eugenia axillaris</i> | Yes | Mesic | <1 | 1-100 | South, Central | South America | BW 74 |
| <i>Eugenia uniflora</i> | Yes | Mesic | <1 | 1-100 | South, Central | South America | |
| <i>Euphorbia alata</i> | | | | | | | |
| <i>Eupatorium capillifolium</i> | | | | | | | |
| <i>Eupatorium leptophyllum</i> | | | | | | | |
| <i>Eupatorium moroni</i> | | | | | | | |
| <i>Eupatorium rotundifolium</i> | | | | | | | |
| <i>Eupatorium serotinum</i> | | | | | | | |
| <i>Euphorbia polystachya</i> | No | Mesic | <1 | 1-100 | South | India, Malay Archipelago, China | BW 59 |
| <i>Euphorbia longan</i> | No | Mesic | <1 | 1-100 | South | India, Malay Archipelago, China | |
| <i>Eustachys glauca</i> | | | | | | | |
| <i>Eustachys petreana</i> | | | | | | | |
| <i>Euthamia caroliniana</i> | | | | | | | |
| <i>Ficus aurea</i> | Yes | Mesic | <1 | 1-100 | South, Central | Old World | BW 73, 580 |
| <i>Ficus microcarpa</i> | Yes | Mesic | <1 | 1-100 | South, Central | Old World | |
| <i>Fimbristylis autumnalis</i> | | | | | | | |
| <i>Fimbristylis cymosa</i> | | | | | | | |
| <i>Fimbristylis dichotoma</i> | | | | | | | |
| <i>Fimbristylis puberula</i> | | | | | | | |
| <i>Fimbristylis schoenoides</i> | No | Mesic | <1 | 10,001-100,000 | Throughout | Tropical Asia | BW 545 |
| <i>Fraxinus caroliniana</i> | | | | | | | |
| <i>Furcraea brevifolia</i> | | | | | | | |
| <i>Furcraea pumila</i> | | | | | | | |
| <i>Furcraea sciropea</i> | | | | | | | |
| <i>Galactia elliptii</i> | | | | | | | |
| <i>Galactia regularis</i> | | | | | | | |
| <i>Galactia volubilis</i> | | | | | | | |
| <i>Galium tritectorium</i> | | | | | | | |
| <i>Gaylussacia dumosa</i> | No | Mesic | <1 | 1-100 | South, Central | Tropical America | BW 56 |
| <i>Gloriosa sp.</i> | No | Mesic | <1 | 1-100 | South, Central | Tropical America | |
| <i>Gnaphalium obtusifolium</i> | | | | | | | |
| <i>Gomphrena serrata</i> | No | Mesic | <1 | 1-100 | Throughout | Tropical America | |
| <i>Gratiola ramosa</i> | | | | | | | |
| <i>Gymnopogon brevifolius</i> | | | | | | | |
| <i>Habenaria floribunda</i> | | | | | | | |
| <i>Hamelia patens</i> | | | | | | | |
| <i>Hedyotis corymbosa</i> | | | | | | | |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, Italicics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|---|------------|---------|---------|--------------|----------------|-------------------|---------|
| <i>Hedysotis procumbens</i> | | | | | | | |
| <i>Hedysotis uniflora</i> | | | | | | | |
| <i>Heleinum amaranthoides</i> | | | | | | | |
| <i>Heleinum pinnatifidum</i> | | | | | | | |
| <i>Heleanthemum corymbosum</i> | | | | | | | |
| <i>Heleanthemum nashii</i> | | | | | | | |
| <i>Heliotropium polystachyon</i> | | | | | | | |
| <i>Heptaptera subaxillans</i> | No | Mesic | <1 | 1-100 | South, Central | Old World Tropics | |
| <i>Hibiscus tiliaceus</i> | No | Mesic | <1 | 1-100 | South, Central | Old World Tropics | |
| <i>Hieracium megacephalon</i> | | | | | | | |
| <i>Hydrocotyle verticillata</i> | Yes | Wet | 1-100 | 1,001-10,001 | Throughout | Old World | BW 85 |
| <i>Hydrocotyle umbellata</i> | | | | | | | |
| <i>Hydrocotyle verticillata</i> | | | | | | | |
| <i>Hypenicum brachyphyllum</i> | | | | | | | |
| <i>Hypenicum cistifolium</i> | | | | | | | |
| <i>Hypenicum fasciculatum</i> | | | | | | | |
| <i>Hypenicum hypericoides</i> | | | | | | | |
| <i>Hypenicum multiflorum</i> | | | | | | | |
| <i>Hypenicum reductum</i> | | | | | | | |
| <i>Hypenicum tetrapetalum</i> | | | | | | | |
| <i>Hypoixis curtissii</i> | | | | | | | |
| <i>Hypoixis juncea</i> | | | | | | | |
| <i>Hypoxis alata</i> | No | Mesic | <1 | 1-100 | South, Central | Tropical America | BW 657 |
| <i>Hypoxis verticillata</i> | No | Mesic | <1 | 1-100 | South, Central | Tropical America | BW 657 |
| <i>Ilex cassine</i> | | | | | | | |
| <i>Ilex glabra</i> | No | Mesic | <1 | 1-100 | Throughout | Old World Tropics | BW 703 |
| <i>Indigofera hirsuta</i> | No | Mesic | <1 | 1-100 | Throughout | Old World Tropics | BW 703 |
| <i>Ipomoea alba</i> | | | | | | | |
| <i>Ipomoea cordatotrioba</i> | | | | | | | |
| <i>Ipomoea indica</i> var. <i>acuminata</i> | | | | | | | |
| <i>Ipomoea sagittata</i> | | | | | | | |
| <i>Iresine diffusa</i> | | | | | | | |
| <i>Itea virginica</i> | | | | | | | |
| <i>Iva microcephala</i> | | | | | | | |
| <i>Juncus dichotomus</i> | | | | | | | |
| <i>Juncus marginatus</i> | | | | | | | |
| <i>Juncus megacephalus</i> | | | | | | | |
| <i>Juncus repens</i> | | | | | | | |
| <i>Juncus scirpoides</i> | | | | | | | |
| <i>Justicia angusta</i> | | | | | | | |
| <i>Kalanchoe pinnata</i> | No | Mesic | <1 | 1-100 | South, Central | Asia | BW 735 |
| <i>Kosteletzkyia virginica</i> | | | | | | | |
| <i>Kyllinga brevifolia</i> | | | | | | | |
| <i>Kyllinga odorata</i> | | | | | | | |
| <i>Lachnanthes caroliniana</i> | | | | | | | |
| <i>Lachnocaulon anceps</i> | | | | | | | |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|--|------------|---------|-----------|--------------|---------------------------|-------------------------------|-------------|
| <i>Lactuca graminifolia</i> | | | | | | | |
| <i>Lagerstroemia indica</i> | No | Mesic | <1 | 1-100 | Throughout | Asia | |
| <i>Lantana camara</i> | Yes | Mesic | <1 | 101-1,000 | Throughout | Tropical America | BW 736 |
| <i>Lechea cernua</i> | | | | | | | |
| <i>Lechea torreyi</i> | | | | | | | |
| <i>Leersia hexandra</i> | | | | | | | |
| <i>Lemma vatividiana</i> | | | | | | | |
| <i>Lepidium virginicum</i> | | | | | | | |
| <i>Liatris chapmanii</i> | | | | | | | |
| <i>Liatris gracilis</i> | | | | | | | |
| <i>Liatris spicata</i> | | | | | | | |
| <i>Liatris tenuifolia</i> | | | | | | | |
| <i>Licania michauxii</i> | | | | | | | |
| <i>Lilium catesbeianum</i> | | | | | | | |
| <i>Linnophila sessiliflora</i> | Yes | Wet | 1-100 | 1,001-10,000 | South, Central, Panhandle | Old World Tropics | BW 106, 275 |
| <i>Linaria canadensis</i> | | | | | | | |
| <i>Lindernia crustacea</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Indo-Malaya | BW 624 |
| <i>Linum floridanum</i> | | | | | | | |
| <i>Lipocarpha aristulata</i> | Yes | Mesic | <1 | 101-1,000 | South, Central | Midwestern and Pacific States | BW 591 |
| <i>Linope muscati</i> | | | | | | | |
| <i>Livistona chinensis</i> | No | Mesic | <1 | 1-100 | South, Central, North | S. Japan, China | |
| <i>Lobelia paludosa</i> | | | | | | | |
| <i>Ludwigia alata</i> | | | | | | | |
| <i>Ludwigia linifolia</i> | | | | | | | |
| <i>Ludwigia maritima</i> | | | | | | | |
| <i>Ludwigia microcarpa</i> | | | | | | | |
| <i>Ludwigia octovalvis</i> | | | | | | | |
| <i>Ludwigia peruviana</i> | | | | | | | |
| <i>Ludwigia repens</i> | | | | | | | |
| <i>Lycopodiella appressa</i> | | | | | | | |
| <i>Lycopodiella cernua</i> | | | | | | | |
| <i>Lycopodium rubellus</i> | | | | | | | |
| <i>Lygodesmia apylia</i> | | | | | | | |
| <i>Lygodium microphyllum</i> | Yes | Wet | 100-1,000 | >100,000 | South, Central | Asia | BW 54, 96 |
| <i>Lyonia fruticosa</i> | | | | | | | |
| <i>Lyonia lucida</i> | | | | | | | |
| <i>Lythrum alatum</i> var. <i>lanceolatum</i> | | | | | | | |
| <i>Magnolia virginiana</i> | | | | | | | |
| <i>Mangifera indica</i> | Yes | Mesic | <1 | 1-100 | South, Central | East Indies | BW 57 |
| <i>Mecardonia acuminata</i> ssp. <i>peninsularis</i> | | | | | | | |
| <i>Metasequoia quinquefolia</i> | Yes | Wet | 1-100 | 1,001-10,000 | South, Central | Australia | BW 118, 595 |
| <i>Melia azedarach</i> | No | Mesic | <1 | 1-100 | Throughout | Asia | BW 727 |
| <i>Micranthemum umbrosum</i> | | | | | | | |
| <i>Mikania cordifolia</i> | | | | | | | |
| <i>Mikania scandens</i> | | | | | | | |
| <i>Mimosa pigra</i> | Yes | Wet | <1 | 1-100 | South, Central | Tropical America | BW 737 |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|---|---------|---------|------------|----------------|-----------------------|------------------|
| <i>Mimosa quadrivalvis</i> | | | | | | |
| <i>Mitreola patiolata</i> | | | | | | |
| <i>Mitrella sessilifolia</i> | | | | | | |
| <i>Momordica charantia</i> | No | Mesic | <1 | 100-1,000 | South, Central, North | BW 713 |
| <i>Monis rubra</i> | | | | | | |
| <i>Muhlenbergia schreberi</i> | | | | | | |
| <i>Murdannia nudiflora</i> | No | Mesic | <1 | 101-1,000 | Throughout | BW 554 |
| <i>Myrica cerifera</i> | | | | | | |
| <i>Nephrolepis cordifolia</i> | Yes | Mesic | <1 | 101-1,000 | Throughout | BW 75 |
| <i>Nephrolepis multiflora</i> | Yes | Mesic | <1 | 101-1,000 | South, Central | BW 114, 617 |
| <i>Nuphar lutea</i> spp. <i>advena</i> | | | | | | |
| <i>Ocotea coriacea</i> | | | | | | |
| <i>Oenothera</i> sp. | | | | | | |
| <i>Ophioglossum palmatum</i> | | | | | | |
| <i>Opismenus hirsutellus</i> | | | | | | |
| <i>Opuntia humifusa</i> | | | | | | |
| <i>Osmunda cinnamomea</i> | | | | | | |
| <i>Osmunda regalis</i> var. <i>spectabilis</i> | | | | | | |
| <i>Oxalis corniculata</i> | | | | | | |
| <i>Oxypolis filiformis</i> | | | | | | |
| <i>Palafoxia feayi</i> | | | | | | |
| <i>Palafoxia integrifolia</i> | | | | | | |
| <i>Panicum dichotomiflorum</i> var. <i>bartowense</i> | | | | | | |
| <i>Panicum hemitomon</i> | | | | | | |
| <i>Panicum hians</i> | | | | | | |
| <i>Panicum maximum</i> | No | Mesic | 1-100 | 101-1,000 | Throughout | Africa |
| <i>Panicum repens</i> | Yes | Wat | 1-100 | 10,001-100,000 | Throughout | Tropical America |
| <i>Panicum rigidulum</i> | | | | | | |
| <i>Panicum tenerum</i> | | | | | | |
| <i>Panicum virgatum</i> | | | | | | |
| <i>Parietaria floridana</i> | | | | | | |
| <i>Pathenocissus quinquefolia</i> | | | | | | |
| <i>Paspalum boscianum</i> | | | | | | |
| <i>Paspalum conjugatum</i> | | | | | | |
| <i>Paspalum distichum</i> | | | | | | |
| <i>Paspalum notatum</i> | No | Mesic | 1-100 | 10,001-100,000 | Throughout | South America |
| <i>Paspalum praecox</i> | | | | | | |
| <i>Paspalum setaceum</i> | | | | | | |
| <i>Paspalum urvillei</i> | No | Mesic | <1 | 101-1,000 | Throughout | South America |
| <i>Passiflora edulis</i> | Yes | Wat | <1 | 1-100 | South, Central | South America |
| <i>Passiflora suberosa</i> | | | | | | |
| <i>Peltandra virginica</i> | | | | | | |
| <i>Pennisetum purpureum</i> | No | Mesic | <1 | 1-100 | Throughout | Africa |
| <i>Persea americana</i> | No | Mesic | <1 | 1-100 | South, Central | Tropical America |
| <i>Persea palustris</i> | | | | | | |
| <i>Phlebodium aureum</i> | | | | | | |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|---|------------|---------|---------|--------------|----------------|------------------------------------|---------|
| <i>Pteridium aquilinum</i> var. <i>caudatum</i> | | | | | | | |
| <i>Pteridium aquilinum</i> var. <i>pseudocaudatum</i> | | | | | | | |
| <i>Pteris vittata</i> | Yes | Mesic | <1 | 101-1,000 | Throughout | Eastern Asia | BW 135 |
| <i>Pterocaulon pyrostachyum</i> | | | | | | | |
| cf. <i>Pteroglossaspis ecristata</i> | | | | | | | |
| <i>Ptilium capillaceum</i> | | | | | | | |
| <i>Quercus chapmanii</i> | | | | | | | |
| <i>Quercus geminata</i> | | | | | | | |
| <i>Quercus laurifolia</i> s.str | | | | | | | |
| <i>Quercus minima</i> | | | | | | | |
| <i>Quercus myrtifolia</i> | | | | | | | |
| <i>Quercus pumila</i> | | | | | | | |
| <i>Quercus virginiana</i> | | | | | | | |
| <i>Rapanea punctata</i> | | | | | | | |
| <i>Rhexia cubensis</i> | | | | | | | |
| <i>Rhexia mariana</i> | | | | | | | |
| <i>Rhexia nuttallii</i> | | | | | | | |
| <i>Rhus copallina</i> | | | | | | | |
| <i>Rhynchosia repens</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | South Africa | |
| <i>Rhynchosia minima</i> | | | | | | | |
| <i>Rhynchospora ciliaris</i> | | | | | | | |
| <i>Rhynchospora colorata</i> | | | | | | | |
| <i>Rhynchospora divergens</i> | | | | | | | |
| <i>Rhynchospora fascicularis</i> | | | | | | | |
| <i>Rhynchospora globularis</i> | | | | | | | |
| <i>Rhynchospora harperi</i> | | | | | | | |
| <i>Rhynchospora intermedia</i> | | | | | | | |
| <i>Rhynchospora inundata</i> | | | | | | | |
| <i>Rhynchospora latifolia</i> | | | | | | | |
| <i>Rhynchospora megatocarpa</i> | | | | | | | |
| <i>Rhynchospora microcarpa</i> | | | | | | | |
| <i>Rhynchospora microcephala</i> | | | | | | | |
| <i>Rhynchospora miliacea</i> | | | | | | | |
| <i>Rhynchospora nitens</i> | | | | | | | |
| <i>Rhynchospora plumosa</i> | | | | | | | |
| <i>Rhynchospora rariflora</i> | | | | | | | |
| <i>Rhynchospora tracyi</i> | | | | | | | |
| <i>Richardia brasiliensis</i> | No | Mesic | <1 | 1-100 | Throughout | South America | BW 61 |
| <i>Richardia grandiflora</i> | No | Mesic | <1 | 101-1,000 | South, Central | South America | BW 639 |
| <i>Richardia scabria</i> | No | Mesic | <1 | 1-100 | Throughout | South America | BW 714 |
| <i>Rivina humilis</i> | | | | | | | |
| <i>Rotala ramosior</i> | | | | | | | |
| <i>Rotboellia cochinchinensis</i> | No | Mesic | <1 | 101-1,000 | Throughout | Tropical Asia | BW 71 |
| <i>Roxstonea regia</i> | Yes | Mesic | <1 | 1-100 | South | Florida, Bahamas, Cuba, Hispaniola | BW 633 |
| <i>Rubus trivialis</i> | | | | | | | |
| <i>Sabal palmetto</i> | | | | | | | |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|--|---------|---------|------------|----------------|-----------------------|------------|
| <i>Sabicea bartramii</i> | | | | | | |
| <i>Sabicea brevifolia</i> | | | | | | |
| <i>Sabicea calycina</i> | | | | | | |
| <i>Saccharum giganteum</i> | | | | | | |
| <i>Sacciolepis indica</i> | Yes | Mesic | 1-100 | >100,000 | Throughout | BW 541 |
| <i>Sacciolepis striata</i> | | | | | | |
| <i>Sagittaria lancifolia</i> | | | | | | |
| <i>Salix caroliniana</i> | | | | | | |
| <i>Salvinia minima</i> | | | | | | |
| <i>Sambucus canadensis</i> | | | | | | |
| <i>Samolus valerandi</i> ssp. <i>pannitifolius</i> | | | | | | |
| <i>Sansevieria hyacinthoides</i> | No | Mesic | <1 | 1-100 | South, Central, North | BW 629 |
| <i>Sarcostemma clausum</i> | | | | | | |
| <i>Sauvagesia cernuus</i> | | | | | | |
| <i>Schefflera actinophylla</i> | Yes | Mesic | <1 | 1-100 | Australia | |
| <i>Schinus terebinthifolius</i> | Yes | Mesic | 100-1,000 | 10,000-100,000 | South, Central, North | BW 43, 596 |
| <i>Schizachyrium stoloniferum</i> | | | | | | |
| <i>Schoenolitton albiflorum</i> | | | | | | |
| <i>Schoenocaulon dubium</i> | | | | | | |
| <i>Schoepfia chrysophylloides</i> | | | | | | |
| <i>Scieria cf. ciliata</i> | | | | | | |
| <i>Scieria georgiana</i> | | | | | | |
| <i>Scieria hirtella</i> | | | | | | |
| <i>Scieria reticularis</i> | | | | | | |
| <i>Scieria triglomerata</i> | | | | | | |
| <i>Scieria verticillata</i> | | | | | | |
| <i>Scoparia dulcis</i> | | | | | | |
| <i>Seneca pendula</i> var. <i>glabrata</i> | Yes | Mesic | <1 | 1-100 | South, Central | |
| <i>Serenoa repens</i> | | | | | | |
| <i>Sesbania vesicaria</i> | | | | | | |
| <i>Setaria parviflora</i> | | | | | | |
| <i>Sida acuta</i> | | | | | | |
| <i>Sida cordifolia</i> | No | Mesic | <1 | 1-100 | Throughout | BW 636 |
| <i>Sida rhombifolia</i> | | | | | | |
| <i>Sideroxylon recinatum</i> | | | | | | |
| <i>Sideroxylon salicifolium</i> | | | | | | |
| <i>Simarouba glauca</i> | | | | | | |
| <i>Sisyrinchium xerophyllum</i> | | | | | | |
| <i>Smilax auriculata</i> | | | | | | |
| <i>Smilax bona-nox</i> | | | | | | |
| <i>Smilax laurifolia</i> | | | | | | |
| <i>Solanum americanum</i> | | | | | | |
| <i>Solanum capsicoides</i> | | | | | | |
| <i>Solanum diphyllum</i> | No | Mesic | <1 | 1-100 | South, Central, North | BW 77, 709 |
| <i>Solidago fistulosa</i> | | | | | | |
| <i>Solidago odora</i> var. <i>chapmanii</i> | | | | | | |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|---|------------|---------|---------|----------------|---------------------------|------------------------|---------|
| Solidago stricta | | | | | | | |
| Solidago latifolia | | | | | | | |
| Sonchus asper | No | Mesic | <1 | 1-100 | Throughout | Europe | |
| Sorghastrum secundum | | | | | | | |
| Sorghum halepense | No | Mesic | <1 | 1-100 | Throughout | Mediterranean | BW 139 |
| Spermatoceps assurgens | | | | | | | |
| Spermatoceps verticillata | No | Mesic | <1 | 1,001-10,000 | South, Central, North | Africa | |
| Spirodela punctata | Yes | Wat | <1 | 10,001-100,000 | Throughout | Old World | BW 66 |
| Sporobolus indicus var. indicus | No | Mesic | <1 | 1,001-10,000 | Throughout | Tropical Asia | BW 142 |
| Sporobolus indicus var. pyramidalis | No | Mesic | 1-100 | 10,001-100,000 | South, Central, Panhandle | Tropical America | BW 718 |
| Sporobolus junceus | | | | | | | BW 44 |
| Stillingia aquatica | | | | | | | |
| Stipulicida setacea | | | | | | | |
| Syagrus romanzoffiana | No | Mesic | <1 | 1-100 | South | S. Brazil to Argentina | GB 1073 |
| Syngonanthus flavidulus | | | | | | | |
| Syngonium podophyllum | Yes | Wat | 1-100 | 1,001-10,000 | South, Central | Central America | BW 626 |
| Syzygium cumini | Yes | Mesic | 1-100 | 101-1,000 | South | Asia | BW 59 |
| Syzygium Jambos | No | Mesic | <1 | 1-100 | South, Central | Asia | BW 72 |
| Taxodium ascendens | | | | | | | |
| Taxodium distichum | | | | | | | |
| Tephrosia rugellii | | | | | | | |
| Thalia geniculata | | | | | | | |
| Thelypteris dentata | Yes | Wat | 1-100 | 101-1,000 | Throughout | Old World Tropics | BW 53 |
| Thelypteris integrifolia | | | | | | | |
| Thelypteris kunthii | | | | | | | |
| Thelypteris semirata | | | | | | | |
| Tiliacaria balbisiana | | | | | | | |
| Tiliacaria fasciculata var. densispica | | | | | | | |
| Tiliacaria paucifolia | | | | | | | |
| Tiliacaria recurvata | | | | | | | |
| Tiliacaria setacea | | | | | | | |
| Tiliacaria usneoides | | | | | | | |
| Tiliacaria utricularia | | | | | | | |
| Tiliacaria variabilis | | | | | | | |
| Toxicodendron radicans | | | | | | | |
| Triadenum virginicum | | | | | | | |
| Trichostema dichotomum | | | | | | | |
| Tridax procumbens | No | Mesic | <1 | 1-100 | South, Central | Tropical America | BW 638 |
| Tridens flavus | | | | | | | |
| Tripsacum dactyloides | | | | | | | |
| Typha domingensis | | | | | | | |
| Urena lobata | | | | | | | |
| Urochloa mutica | No | Wat | 1-100 | 101-1,000 | South, Central | Africa | BW 553 |
| Urochloa ramosa | No | Mesic | <1 | 1-100 | Throughout | Asia | BW 622 |
| Urticularia cornuta | | | | | | | |

Loxahatchee River Corridor - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size. | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|--------------|----------------|------------------------|---------|
| <i>Utricularia foliosa</i> | | | | | | | |
| <i>Utricularia gibba</i> | | | | | | | |
| <i>Utricularia purpurea</i> | | | | | | | |
| <i>Utricularia simulans</i> | | | | | | | |
| <i>Utricularia subulata</i> | | | | | | | |
| <i>Vaccinium myrsinites</i> | | | | | | | |
| <i>Vaccinium stamineum</i> | | | | | | | |
| <i>Verbena virginica</i> | | | | | | | |
| <i>Vernonia cinercea</i> | No | Mesic | <1 | 101-1,000 | South, Central | Tropical America | BW 137 |
| <i>Vicia acutifolia</i> | | | | | | | |
| <i>Vigna luteola</i> | | | | | | | |
| <i>Vitis cinerea</i> var. <i>floridana</i> | | | | | | | |
| <i>Vitis rotundifolia</i> | | | | | | | |
| <i>Vitis shuttleworthii</i> | | | | | | | |
| <i>Vitaria lineata</i> | | | | | | | |
| <i>Wedelia trilobata</i> | No | Mesic | <1 | 101-1,000 | Throughout | West Indies | |
| <i>Wolffia cf. columbiana</i> | | | | | | | |
| <i>Woodwardia virginica</i> | | | | | | | |
| <i>Xanthosoma sagittifolium</i> | No | Mesic | <1 | 1-100 | South, Central | Tropical America | BW 627 |
| <i>Ximenia americana</i> | | | | | | | |
| <i>Xyris ambigua</i> | | | | | | | |
| <i>Xyris brevifolia</i> | | | | | | | |
| <i>Xyris caroliniana</i> | | | | | | | |
| <i>Xyris difformis</i> var. <i>floridana</i> | | | | | | | |
| <i>Xyris elliottii</i> | | | | | | | |
| <i>Xyris flabelliformis</i> | | | | | | | |
| <i>Xyris Jupical</i> | Yes | Mesic | 1-100 | 1,001-10,000 | Throughout | Tropical America | BW 609 |
| <i>Xyris smalliana</i> | | | | | | | |
| <i>Youngia japonica</i> | Yes | Mesic | <1 | 1,001-10,000 | Throughout | SE Asia | BW 631 |
| <i>Yucca aloifolia</i> | No | Mesic | <1 | 1-100 | Throughout | West Indies and Mexico | |

Loxahatchee River Corridor - Endangered and Threatened Taxa

| | State Status | Habitat | Pop. Size. | # Stations | Status | Natural Range | Voucher |
|---|---------------------|----------------|-------------------|-------------------|---------------|----------------------|---|
| <i>Campyloneurum phyllitidis</i> | E | Wet | <1 | 1-100 | Throughout | Stable | Florida to South America BW 116 |
| <i>Chrysophyllum oliviforme</i> | E | Mesic | <1 | 1-100 | | 1 Stable | Florida, Belize, Greater Antilles |
| <i>Eulophia alta</i> | T | Wet | <1 | 101-1,000 | Throughout | Stable | Florida, West Indies, Central and South America |
| <i>Lechea cernua</i> | T | Xeric | <1 | 1-100 | | 1 Stable | Florida |
| <i>Lilium catesbaei</i> | T | Mesic | <1 | 1-100 | | 2 Stable | Va. to La. and FL. |
| <i>Ophioglossum palmatum</i> | E | Mesic | <1 | 101-1,000 | 1 (scattered) | Decreasing | South Fla., tropical America, and Old World BW 564 |
| <i>Polygala smallii</i> | E | Mesic | <1 | 1-100 | | 4 Stable | South and central Florida BW 140 |
| <i>Thelypteris serrata</i> | E | Wet | 1-100 | 1,001-10,000 | Throughout | Stable? | Florida to South America BW 261, 262 |
| <i>Tillandsia balbisiana</i> | T | Wet | 1-100 | 101-1,000 | Throughout | Stable | Florida to South America BW 52 |
| <i>Tillandsia fasciculata var. <i>densipila</i></i> | E | Mesic | 1-100 | 101-1,000 | Throughout | Stable | Florida to Central America |
| <i>Tillandsia utriculata</i> | E | Wet | <1 | 101-1,000 | Throughout | Stable | Florida to South America BW 706 |
| <i>Tillandsia variabilis</i> | T | Wet | <1 | 101-1,000 | Throughout | Stable | Florida to South America GB 1070 |

Nicodemus Slough - Preliminary List of Vascular Plants

Bold = Exotic, /Native = Cultivated

| Species | Nat./Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|---------|--------------|-----------------------|------------------|---------|
| <i>Acer rubrum</i> | | | | | | | |
| <i>Aceria oppositifolia</i> var. <i>repens</i> | | | | | | | |
| <i>Aeschynomene americana</i> | | | | | | | |
| <i>Aldrovanda philoxeroides</i> | Yes | Wet | 1-100 | 1,001-10,000 | Throughout | South America | |
| <i>Amaranthus australis</i> | No | Mesic | <1 | 101-1,000 | Throughout | India | |
| <i>Amaranthus spinosus</i> | | | | | | | |
| <i>Ambrosia artemisiifolia</i> | | | | | | | |
| <i>Ammannia latifolia</i> | | | | | | | |
| <i>Ampelopsis arborea</i> | | | | | | | |
| <i>Andropogon glomeratus</i> var. <i>hirsutior</i> | | | | | | | |
| <i>Andropogon glomeratus</i> var. <i>pumilus</i> | | | | | | | |
| <i>Andropogon gyrans</i> var. <i>gyrans</i> | | | | | | | |
| <i>Andropogon virginicus</i> var. <i>virginicus</i> | | | | | | | |
| <i>Ardisia pallida</i> | No | Mesic | <1 | 101-1,000 | South, Central | South America | |
| <i>Asclepias curassavica</i> | | | | | | | |
| <i>Asclepias incarnata</i> | | | | | | | |
| <i>Aster subulatus</i> | | | | | | | |
| <i>Axonopus compressus</i> | | | | | | | |
| <i>Axonopus furcatus</i> | | | | | | | |
| <i>Azolla caroliniana</i> | | | | | | | |
| <i>Baccharis glomeruliflora</i> | | | | | | | |
| <i>Baccharis halimifolia</i> | | | | | | | |
| <i>Bacopa caroliniana</i> | | | | | | | |
| <i>Bacopa monnieri</i> | | | | | | | |
| <i>Bidens alba</i> var. <i>radiata</i> | | | | | | | |
| <i>Blechnum serrulatum</i> | | | | | | | |
| <i>Blechnum pyramidatum</i> | No | Mesic | <1 | 101-1,000 | South, Central, North | Tropical America | |
| <i>Boehmeria cylindrica</i> | | | | | | | |
| <i>Bucinia americana</i> | | | | | | | |
| <i>Calliprora americana</i> | | | | | | | |
| <i>Caystegia sepium</i> ssp. <i>limnophila</i> | | | | | | | |
| <i>Canna flaccida</i> | | | | | | | |
| <i>Carex vexans</i> | | | | | | | |
| <i>Ceratodon purpureus</i> | | | | | | | |
| <i>Ceratopteris thalictroides</i> | | | | | | | |
| <i>Cenchrus incertus</i> | | | | | | | |
| <i>Centella asiatica</i> | | | | | | | |
| <i>Cephaelanthus occidentalis</i> | | | | | | | |
| <i>Chamaescista fasciculata</i> | | | | | | | |
| <i>Chamaecrista nictitans</i> var. <i>aspera</i> | | | | | | | |
| <i>Chamaesyce blodgettii</i> | | | | | | | |
| <i>Chamaesyce cornifolia</i> | | | | | | | |
| <i>Chamaesyce hirta</i> | | | | | | | |
| <i>Chamaesyce hypericifolia</i> | | | | | | | |
| <i>Chamaesyce hyssopifolia</i> | | | | | | | |
| <i>Chenopodium ambrosioides</i> | No | Mesic | 1-100 | 1,001-10,000 | Throughout | Tropical America | |
| <i>Cirsium horridulum</i> | | | | | | | |
| <i>Cladium laevigatum</i> | | | | | | | |
| <i>Coleorachis rugosa</i> | | | | | | | |
| <i>Commelinina diffusa</i> | | | | | | | |
| <i>Conoclinium coelestinum</i> | | | | | | | |
| <i>Conyzia canadensis</i> var. <i>pusilla</i> | | | | | | | |
| <i>Crocosmia leavenworthii</i> | | | | | | | |
| <i>Crotalaria lanceolata</i> | No | Mesic | <1 | 101-1,000 | Throughout | Africa | BW 455 |
| <i>Crotalaria rotundifolia</i> | | | | | | | |

Nicodemus Slough - Preliminary List of Vascular Plants

Bold = Exotic, **Italics** = Cultivated

| Species | Nat. Area? | Habitat | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|-----------|----------------|----------------------------|-------------------|
| <i>Croton glandulosus</i> var. <i>glandulosus</i> | Yes | Wet | 1-100 | 1,001-10,000 | Throughout | |
| <i>Cuphea carthagenensis</i> | | | | | South and Central America? | BW 200 |
| <i>Cuscuta cf. indecora</i> | | | | | | |
| <i>Cyananthum angustifolium</i> | | | | | | |
| <i>Cynanchum scoparium</i> | | | | | | |
| <i>Cynodon dactylon</i> | No | Mesic | 1-100 | 1,001-10,000 | Throughout | |
| <i>Cyperus compressus</i> | | | | | | BW 472 |
| <i>Cyperus croceus</i> | | | | | | |
| <i>Cyperus distinctus</i> | | | | | | |
| <i>Cyperus flavescens</i> | | | | | | |
| <i>Cyperus haspan</i> | | | | | | |
| <i>Cyperus odoratus</i> | | | | | | |
| <i>Cyperus polystachyos</i> | No | Wet | 1-100 | 1-100 | South, Central Africa | |
| <i>Cyperus prolifer</i> | | | | | | |
| <i>Cyperus retrofractus</i> | | | | | | |
| <i>Cyperus rotundus</i> | No | Mesic | 1-100 | 1,001-10,000 | Throughout | |
| <i>Cyperus surinamensis</i> | | | | | | Eurasia |
| <i>Dactyloctenium aegyptium</i> | No | Mesic | 1-100 | 10,001-100,000 | Throughout | |
| <i>Desmodium incanum</i> | | | | | | Old World Tropics |
| <i>Desmodium trifolium</i> | No | Mesic | <1 | 101-1,000 | Old World | BW 206 |
| <i>Dichanthelium commutatum</i> | | | | | | |
| <i>Dichanthelium dichotomum</i> | | | | | | |
| <i>Dichanthelium encyclioides</i> | | | | | | |
| <i>Dicliptera sexangulata</i> | | | | | | |
| <i>Diiodia virginiana</i> | | | | | | |
| <i>Diospyros virginiana</i> | | | | | | |
| <i>Drimaria condita</i> | | | | | | |
| <i>Echinochloa paludigena</i> | | | | | | |
| <i>Echinochloa walteri</i> | | | | | | |
| <i>Eclipta umbellata</i> | | | | | | |
| <i>Ectrosia prostrata</i> | | | | | | |
| <i>Eleocharis baldwinii</i> | | | | | | |
| <i>Eleocharis elongata</i> | | | | | | |
| <i>Eleocharis intensa</i> | | | | | | |
| <i>Erechtites hieracifolia</i> | | | | | | |
| <i>Eryngium baldwinii</i> | | | | | | |
| <i>Euphorbia alta</i> | | | | | | |
| <i>Eupatorium capillifolium</i> | | | | | | |
| <i>Eupatorium leptophyllum</i> | | | | | | |
| <i>Eustachys glauca</i> | | | | | | |
| <i>Eustachys petraea</i> | | | | | | |
| <i>Euthamia caroliniana</i> | | | | | | |
| <i>Ficus aurea</i> | | | | | | |
| <i>Fimbristylis autumnalis</i> | | | | | | |
| <i>Fimbristylis caroliniana</i> | | | | | | |
| <i>Fimbristylis cymosa</i> | | | | | | |
| <i>Fimbristylis puberula</i> | | | | | | |
| <i>Fraxinus caroliniana</i> | | | | | | |
| <i>Fuirena breviseta</i> | | | | | | |
| <i>Fuirena pumila</i> | | | | | | |
| <i>Gallium theodorum</i> | | | | | | |
| <i>Gaura angustifolia</i> | | | | | | |
| <i>Heterotheca subaxillaris</i> | | | | | | |

Nicodemus Slough - Preliminary List of Vascular Plants

Bold = Exotic; Italics = Cultivated

| Species | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|-----------|--------------|-----------------------|-------------------|---------------|
| Hibiscus grandiflorus | | | | | | | |
| Hydrobia verticillata | Yes | Wet | 1-100 | 1,001-10,000 | Throughout | Old World | |
| Hydrocole umbellata | | | | | | | |
| Hydrocole verticillata | | | | | | | |
| Hydrodea corymbosa | | | | | | | |
| Hymenachne amplexicaulis | Yes | Wet | 1-100 | 101-1,000 | South, Central | West Indies | BW 485 |
| Hypericum hypericoides | | | | | | | |
| Hypoxis alata | | | | | | | |
| Hypoxis verticillata | Yes | Mesic | 1-100 | 1,001-10,000 | South, Central | Tropical America | |
| Iresine diffusa | | | | | | | |
| Juncus marginatus | | | | | | | |
| Juncus megacephalus | | | | | | | |
| Juniperus virginiana | | | | | | | |
| Kosteletzkyia virginica | | | | | | | |
| Lachnanthes caroliniana | | | | | | | |
| Leersia hexandra | | | | | | | |
| Lepidium virginicum | | | | | | | |
| Lindernia grandiflora | | | | | | | |
| Lipocarpha micrantha | | | | | | | |
| Ludwigia maritima | | | | | | | |
| Ludwigia octovalvis | | | | | | | |
| Ludwigia peruviana | | | | | | | |
| Ludwigia repens | | | | | | | |
| Luzula fluitans | | | | | | | |
| Lythrum alatum var. lanceolatum | No | Mesic | <1 | 101-1,000 | South, Central | Panhandle | BW 189 |
| Macropollium latifoloides | | | | | | | |
| Melaleuca quinquenervia | Yes | Wet | 1-100 | 1,000-10,000 | South, Central | Australia | |
| Melothria pendula | | | | | | | |
| Mikania scandens | | | | | | | |
| Mitroeda petiolarata | | | | | | | |
| Momordica charantia | Yes | Mesic | <1 | 101-1,000 | South, Central, North | Old World Tropics | |
| Muhlenbergia schreberi | | | | | | | |
| Myrica cerifera | | | | | | | |
| Naia quadrangularis | | | | | | | |
| Nephrolepis sp. | | | | | | | |
| Nuphar lutea ssp. advena | | | | | | | |
| Osmunda regalis var. spectabilis | | | | | | | |
| Oxalis corniculata | | | | | | | |
| Panicum dichotomiflorum var. bartowense | | | | | | | |
| Panicum hemitomon | | | | | | | |
| Panicum hiemis | | | | | | | |
| Panicum repens | Yes | Wet | 1-100 | 1,001-10,000 | Throughout | Tropical America | BW 435 |
| Panicum rigidulum | | | | | | | |
| Pathenocissus quinquefolia | | | | | | | |
| Paspalidium germinatum | | | | | | | |
| Paspalum conjugatum | | | | | | | |
| Paspalum floridanum | | | | | | | |
| Paspalum notatum | Yes | Mesic | 101-1,000 | >100,000 | Throughout | South America | BW 474 |
| Paspalum praecox | | | | | | | |
| Paspalum repens | | | | | | | |
| Paspalum urvillei | No | Mesic | <1 | 1,001-10,000 | Throughout | South America | BW 433 |
| Pectis prostrata | | | | | | | |
| Phleodium aureum | | | | | | | |

Nicodemus Slough - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| Species | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|-----------|----------------|-----------------------|---------------------|---------|
| <i>Phyla nodiflora</i> | | | | | | | |
| <i>Physalis angulata</i> | | | | | | | |
| <i>Physalis walleri</i> | | | | | | | |
| <i>Phytolacca americana</i> | | | | | | | |
| <i>Pitavia striatula</i> | Yes | Wet | 1-100 | 101-1,000 | Throughout | Tropical America | BW 810 |
| <i>Pitcairnia odorata</i> | | | | | | | |
| <i>Pitchea rosea</i> | | | | | | | |
| <i>Poinsettia cyathophora</i> | | | | | | | |
| <i>Poinsettia heterophylla</i> | | | | | | | |
| <i>Polygonia grandiflora</i> | | | | | | | |
| <i>Polygonum densiflorum</i> | | | | | | | |
| <i>Polygonum hydropiperoides</i> | | | | | | | |
| <i>Polygonum punctatum</i> | | | | | | | |
| <i>Polygonum setaceum</i> | | | | | | | |
| <i>Polyodium polypodioides</i> var. <i>richauxianum</i> | | | | | | | |
| <i>Polypteron procumbens</i> | | | | | | | |
| <i>Pontederia cordata</i> | | | | | | | |
| <i>Pontiaca oloracea</i> | No | Mesic | <1 | 101-1,000 | Throughout | Old World | |
| <i>Potentilla pilosa</i> | | | | | | | |
| <i>Prosopisnaca palustris</i> | | | | | | | |
| <i>Pteris vittata</i> | Yes | Mesic | <1 | 101-1,000 | Throughout | Eastern Asia | |
| <i>Ptilimnium capillaceum</i> | | | | | | | |
| <i>Quercus virginiana</i> | | | | | | | |
| <i>Rhexia mariana</i> | | | | | | | |
| <i>Rhus copallina</i> | | | | | | | |
| <i>Rhynchospora colorata</i> | | | | | | | |
| <i>Rhynchospora divergens</i> | | | | | | | |
| <i>Rhynchospora fascicularis</i> | | | | | | | |
| <i>Rhynchospora inundata</i> | | | | | | | |
| <i>Rhynchospora nitricarpa</i> | | | | | | | |
| <i>Rhynchospora nitens</i> | | | | | | | |
| <i>Rhynchospora odorata</i> | | | | | | | |
| <i>Ricina humilis</i> | | | | | | | |
| <i>Rotala ramosior</i> | | | | | | | |
| <i>Rubus trivialis</i> | | | | | | | |
| <i>Sabal palmetto</i> | | | | | | | |
| <i>Saccharum giganteum</i> | | | | | | | |
| <i>Saccobolus stictata</i> | | | | | | | |
| <i>Sagittaria filiformis</i> | | | | | | | |
| <i>Sagittaria lancifolia</i> | | | | | | | |
| <i>Sagittaria latifolia</i> | | | | | | | |
| <i>Salix caroliniana</i> | | | | | | | |
| <i>Saxifraga mertensiana</i> | | | | | | | |
| <i>Saxifraga minima</i> | | | | | | | |
| <i>Samolus valerandi</i> ssp. <i>parviflorus</i> | | | | | | | |
| <i>Saururus cernuus</i> | | | | | | | |
| <i>Schizandra terebinthifolia</i> | Yes | Mesic | 100-1,000 | 10,001-100,000 | South, Central, North | Brazil and Paraguay | |
| <i>Schipus purgans</i> | | | | | | | |
| <i>Scirpus latifoliusmontani</i> | | | | | | | |
| <i>Sciaria reticularis</i> | | | | | | | |
| <i>Scoparia dulcis</i> | | | | | | | |
| <i>Senna obtusifolia</i> | No | Mesic | <1 | 1-100 | Throughout | Origin Uncertain | |
| <i>Sesbania herbacea</i> | | | | | | | |

Nicodemus Slough - Preliminary List of Vascular Plants

Bold = Exotic, *italics* = Cultivated

| Species | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|----------------|---------------------------|----------------------|---------|
| <i>Setaria parviflora</i> | | | | | | | |
| <i>Sida acuta</i> | | | | | | | |
| <i>Sida rhombifolia</i> | | | | | | | |
| <i>Sideroxylon declinatum</i> spp. <i>austrofloridense</i> | | | | | | | |
| <i>Smilax australis</i> | | | | | | | |
| <i>Smilax bona-nox</i> | | | | | | | |
| <i>Smilax tamnoides</i> | | | | | | | |
| <i>Solanum americanum</i> | | | | | | | |
| <i>Solanum viarum</i> | Yes | Mesic | <1 | 1-100 | Throughout | Argentina and Brazil | BW 822 |
| <i>Solidago gigantea</i> | | | | | | | |
| <i>Solidago sempervirens</i> | | | | | | | |
| <i>Sorghum halepense</i> | No | Mesic | <1 | 1-100 | Throughout | Mediterranean region | BW 479 |
| <i>Spermacoce assurgens</i> | | | | | | | |
| <i>Sporodesmia punctata</i> | No | Wet | <1 | 10,001-100,000 | Throughout | Old World | BW 811 |
| <i>Sporobolus indicus</i> var. <i>Indicus</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Tropical Asia | BW 473 |
| <i>Sporobolus indicus</i> var. <i>pyramidalis</i> | No | Mesic | 1-100 | 10,001-100,000 | South, Central, Panhandle | Tropical America | BW 454 |
| <i>Stenophyllum secundatum</i> | | | | | | | |
| <i>Tecoma distichum</i> | | | | | | | |
| <i>Tecophilium canadense</i> | | | | | | | |
| <i>Thelypteris intermedia</i> | | | | | | | |
| <i>Thelypteris turneri</i> | | | | | | | |
| <i>Thlaspidia bahiensana</i> | | | | | | | |
| <i>Thlaspidia recurvata</i> | | | | | | | |
| <i>Thlaspidia setacea</i> | | | | | | | |
| <i>Thlaspidia usneoides</i> | | | | | | | |
| <i>Thlaspidia utricularia</i> | | | | | | | |
| <i>Toxicodendron radicans</i> | | | | | | | |
| <i>Typea dominicensis</i> | | | | | | | |
| <i>Typha latifolia</i> | | | | | | | |
| <i>Urena lobata</i> | | | | | | | |
| <i>Urochloa mutica</i> | Yes | Wet | 1-100 | 101-1,000 | South, Central | Africa | BW 814 |
| <i>Utricularia gibba</i> | | | | | | | |
| <i>Utricularia foliosa</i> | | | | | | | |
| <i>Verbena scabra</i> | | | | | | | |
| <i>Verbesina virginica</i> | | | | | | | |
| <i>Vicia acutifolia</i> | | | | | | | |
| <i>Vigna luteola</i> | | | | | | | |
| <i>Vitis cinerea</i> var. <i>Iorkiana</i> | | | | | | | |
| <i>Vitis rotundifolia</i> | | | | | | | |
| <i>Vittaria lineata</i> | | | | | | | |
| <i>Woodwardia virginica</i> | | | | | | | |
| <i>Xyris jupicai</i> | Yes | Mesic | 1-100 | 101-1,000 | Throughout | Tropical America | |

Nicodemus Slough - Endangered and Threatened Taxa

| | State | Status | Habitat | Acreage | Pop. Size | # Stations | Status | Natural Range | Voucher |
|------------------------------|-------|--------|---------|-------------|------------|------------|---|--|---------|
| <i>Eulophia alta</i> | T | Wet | 1-100 | 1001-10,000 | Throughout | Stable | Florida, West Indies, Central and South America | | |
| <i>Tillandsia balbisiana</i> | T | Mesic | <1 | 1-100 | | 1 | Stable | Florida to South America | |
| <i>Tillandsia utriculata</i> | E | Mesic | <1 | 101-1,000 | | 2 | Stable | Georgia and Florida, West Indies, Mexico, Northern South America, Venezuela. | BW 823 |

Pal-Mar - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|-----------|----------------|---------------|---------|
| <i>Acer rubrum</i> | | | | | | | |
| <i>Acrostichum danaeifolium</i> | | | | | | | |
| <i>Agalinis fasciculata</i> | | | | | | | |
| <i>Agalinis linifolia</i> | | | | | | | |
| <i>Ailanthus lebbeck</i> | No | Mesic | <1 | 1-100 | South, Central | Asia | |
| <i>Aletris lutea</i> | | | | | | | |
| <i>Alseis carpus ovalifolius</i> | No | Mesic | <1 | 101-1,000 | Throughout | Old World | BW 663 |
| <i>Amelanchier alnifolia</i> | | | | | | | |
| <i>Amphicarpum muellerbergianum</i> | | | | | | | |
| <i>Andropogon glomeratus</i> var. <i>glaucoensis</i> | | | | | | | |
| <i>Andropogon glomeratus</i> var. <i>hirsutior</i> | | | | | | | |
| <i>Andropogon glomeratus</i> var. <i>pumilus</i> | | | | | | | |
| <i>Andropogon gyrans</i> var. <i>gyrans</i> | | | | | | | |
| <i>Andropogon ternarius</i> | | | | | | | |
| <i>Andropogon virginicus</i> var. <i>decipiens</i> | | | | | | | |
| <i>Andropogon virginicus</i> var. <i>glaucus</i> | | | | | | | |
| <i>Annona glabra</i> | | | | | | | |
| <i>Aristida beyrichiana</i> | | | | | | | |
| <i>Aristida palustris</i> | | | | | | | |
| <i>Aristida patula</i> | | | | | | | |
| <i>Aristida purpureocans</i> | | | | | | | |
| <i>Aristida frizomophora</i> | | | | | | | |
| <i>Aristida spiciformis</i> | | | | | | | |
| <i>Asclepias lanceolata</i> | | | | | | | |
| <i>Asimina reticulata</i> | | | | | | | |
| <i>Aster dumosus</i> | | | | | | | |
| <i>Axonopus fissifolius</i> | | | | | | | |
| <i>Axonopus furcatus</i> | | | | | | | |
| <i>Baccharis glomeruliflora</i> | | | | | | | |
| <i>Bacopa caroliniana</i> | | | | | | | |
| <i>Bacopa monnierii</i> | | | | | | | |
| <i>Bartsia paniculata</i> | | | | | | | |
| <i>Bartsia virginica</i> | | | | | | | |
| <i>Betania racemosa</i> | | | | | | | |
| <i>Bidens alba</i> var. <i>radiata</i> | | | | | | | |
| <i>Bignonia ruddiae</i> ssp. <i>australis</i> | | | | | | | |
| <i>Blechnum semialatum</i> | | | | | | | |
| <i>Boehmeria cylindrica</i> | | | | | | | |
| <i>Buchnera americana</i> | | | | | | | |
| <i>Bucida cf. buceras</i> | No | Mesic | <1 | 1-100 | South | | GB 1063 |
| <i>Burmannia capitata</i> | | | | | | | |
| <i>Carphophorus camosus</i> | | | | | | | |
| <i>Carphophorus compositus</i> | | | | | | | |
| <i>Cassytha filiformis</i> | No | Mesic | <1 | 101-1,000 | South, Central | Australia | BW 238 |
| <i>Casuarina glauca</i> | | | | | | | |
| <i>Cenchrus incertus</i> | | | | | | | |
| <i>Centella asiatica</i> | | | | | | | |
| <i>Centrosema virginianum</i> | | | | | | | |

Pal-Mar - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|---------|--------------|---------------|----------------------------|---------|
| <i>Cephaelanthus occidentalis</i> | | | | | | | |
| <i>Chamaecrista fasciculata</i> | | | | | | | |
| <i>Chamaecrista nictitans</i> var. <i>aspera</i> | | | | | | | |
| <i>Chamaesycea blodgettii</i> | | | | | | | |
| <i>Chamaesycea hypericifolia</i> | | | | | | | |
| <i>Chamisia tomentosa</i> | | | | | | | |
| <i>Chrysobalanus icaco</i> | | | | | | | |
| <i>Cirsium horridulum</i> | | | | | | | |
| <i>Cladium famalicense</i> | | | | | | | |
| <i>Clematis baldwinii</i> | | | | | | | |
| <i>Colorachis rugosa</i> | | | | | | | |
| <i>Comidendron coelostitnum</i> | | | | | | | |
| <i>Conyza canadensis</i> var. <i>pusilla</i> | | | | | | | |
| <i>Coreopsis floridana</i> | | | | | | | |
| <i>Coreopsis leavenworthii</i> | | | | | | | |
| <i>Crotalaria lanceolata</i> | No | Mesic | <1 | 101-1,000 | Throughout | Africa | |
| <i>Crotalaria pallida</i> var. <i>obovata</i> | No | Mesic | <1 | 101-1,000 | Throughout | Africa | |
| <i>Crotalaria rotundifolia</i> | | | | | | | |
| <i>Crotalaria spectabilis</i> | No | Mesic | <1 | 101-1,000 | Throughout | Old World Tropics | |
| <i>Croton glandulosus</i> var. <i>glandulosus</i> | | | | | | | |
| <i>Clemum aromaticum</i> | No | Mesic | <1 | 1-100 | Throughout | Old World Tropics | |
| <i>Cucumis melo</i> | Yes | Wet | 1-100 | 101-1,000 | Throughout | South and Central America? | |
| <i>Cuphea carthagenensis</i> | | | | | | | |
| <i>Cyperus croceus</i> | | | | | | | |
| <i>Cyperus flavescens</i> | | | | | | | |
| <i>Cyperus haspan</i> | | | | | | | |
| <i>Cyperus ligularis</i> | | | | | | | |
| <i>Cyperus polystachyos</i> | | | | | | | |
| <i>Cyperus retrorsus</i> | | | | | | | |
| <i>Cyperus surinamensis</i> | No | Mesic | 1-100 | 1,001-10,000 | Throughout | Old World Tropics | |
| <i>Dactyloctenium aegyptium</i> | | | | | | | |
| <i>Desmodium incanum</i> | | | | | | | |
| <i>Desmodium triflorum</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Old World | |
| <i>Dichanthelium aciculare</i> | | | | | | | |
| <i>Dichanthelium dichotomum</i> | | | | | | | |
| <i>Dichanthelium ensifolium</i> | | | | | | | |
| <i>Dichanthelium erectifolium</i> | | | | | | | |
| <i>Dichanthelium leucothrix</i> | | | | | | | |
| <i>Dichanthelium portoricense</i> | | | | | | | |
| <i>Dichanthelium scabiosculum</i> | | | | | | | |
| <i>Dodonaea angustifolia</i> | | | | | | | |
| <i>Drosera capillaris</i> | | | | | | | |
| <i>Dyschoriste angusta</i> | | | | | | | |
| <i>Echinocloa paludigena</i> | | | | | | | |
| <i>Eleocharis baldwinii</i> | | | | | | | |
| <i>Eleocharis cellulosa</i> | | | | | | | |
| <i>Eleocharis elongata</i> | | | | | | | |
| <i>Eleocharis geniculata</i> | | | | | | | |

Pai-Mar - Preliminary List of Vascular Plants

Bold = Exotic, Italicics = Cultivated

| Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|------------|---------------------------------|---------|-----------|----------------|-----------------------|-------------------|
| | <i>Eleocharis interstincta</i> | | | | | |
| | <i>Elephantopus elatus</i> | | | | | |
| No | <i>Emilia sonchifolia</i> | Mesic | <1 | 101-1,000 | South, Central, North | |
| | <i>Encyclia tampensis</i> | | | | Old World | |
| No | <i>Eragrostis amabilis</i> | Mesic | <1 | 1-100 | Throughout | |
| | <i>Eragrostis atrovirens</i> | Wet | 1-100 | 1,001-10,000 | Southeast Asia | BW 665 GB 1030 |
| | <i>Eragrostis elliotii</i> | | | | | |
| | <i>Eragrostis spectabilis</i> | | | | | |
| | <i>Eragrostis virginica</i> | | | | | |
| No | <i>Eremochloa opaciflora</i> | Mesic | 1-100 | >100,000 | Throughout | |
| | <i>Eriogonum quercifolius</i> | | | | Southeast Asia | BW 662 |
| | <i>Eriogonum venustum</i> | | | | | |
| | <i>Ernocalyx decangulare</i> | | | | | |
| | <i>Ernocalyx ravenelli</i> | | | | | |
| | <i>Eryngium yuccifolium</i> | | | | | |
| | <i>Euphorbia alta</i> | | | | | |
| | <i>Eupatorium capillifolium</i> | | | | | |
| | <i>Eupatorium leptophyllum</i> | | | | | |
| | <i>Eupatorium mikanioides</i> | | | | | |
| | <i>Eupatorium mohrii</i> | | | | | |
| | <i>Eustachys glauca</i> | | | | | |
| | <i>Eustachys petraea</i> | | | | | |
| | <i>Euthamia caroliniana</i> | | | | | |
| | <i>Evhovulus sanctus</i> | | | | | |
| | <i>Ficus aurea</i> | | | | | |
| | <i>Fimbristylis autumnalis</i> | | | | | |
| | <i>Fimbristylis cymosa</i> | | | | | |
| | <i>Flaveria linearis</i> | | | | | |
| | <i>Fultrena pumila</i> | | | | | |
| | <i>Fuliginea scirpoidea</i> | | | | | |
| | <i>Galecia elliptii</i> | | | | | |
| | <i>Galecia volubilis</i> | | | | | |
| | <i>Gaura angustifolia</i> | | | | | |
| | <i>Gordonia lasianthus</i> | | | | | |
| | <i>Gratiola pilosa</i> | | | | | |
| | <i>Habenaria floribunda</i> | | | | | |
| | <i>Hebdomis uniflora</i> | | | | | |
| | <i>Helenium pinnatifidum</i> | | | | | |
| | <i>Heliotropium polypodium</i> | | | | | |
| | <i>Heterotheca subaxillaris</i> | | | | | |
| | <i>Hydrocotyle sp.</i> | | | | | |
| | <i>Hymenocallis palmeri</i> | No | 1-100 | 10,001-100,000 | South, Central, North | BW 240 |
| | <i>Hypanthocleia rufa</i> | | | | Africa | |
| | <i>Hypericum brachyphyllum</i> | | | | | |
| | <i>Hypericum diffusum</i> | | | | | |
| | <i>Hypericum fasciculatum</i> | | | | | |
| | <i>Hypericum hypericoides</i> | | | | | |
| | <i>Hypericum tetrapetalum</i> | | | | | |
| | <i>Hypoxis juncea</i> | | | | | |

Pai-Mar - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|----------------|-----------------------|--|---------|
| <i>Hyptis alata</i> | | | | | | | |
| <i>Ilex cassine</i> | | | | | | | |
| <i>Ilex glabra</i> | | | | | | | |
| <i>Imperata cylindrica</i> | No | Mesic | <1 | 101-1,000 | South, Central, North | Mediterranean to South Africa, India, southeast Asia to Australia. | BW 239 |
| <i>Ipomoea sagittata</i> | | | | | | | |
| <i>Iresine diffusa</i> | | | | | | | |
| <i>Iva microcephala</i> | | | | | | | |
| <i>Juncus marginatus</i> | | | | | | | |
| <i>Juncus megacephalus</i> | | | | | | | |
| <i>Juncus scirpoides</i> | | | | | | | |
| <i>Lachnanthes caroliniana</i> | | | | | | | |
| <i>Lachnocaulon articulatum</i> | No | Mesic | <1 | 101-1,000 | Throughout | Tropical America | |
| <i>Lamium camara</i> | | | | | | | |
| <i>Leersia hexandra</i> | | | | | | | |
| <i>Liatris gracilis</i> | | | | | | | |
| <i>Liatris spicata</i> | | | | | | | |
| <i>Lilium catesbaei</i> | | | | | | | |
| <i>Linum medium</i> var. <i>texanum</i> | | | | | | | |
| <i>Lobelia glandulosa</i> | | | | | | | |
| <i>Ludwigia alata</i> | | | | | | | |
| <i>Ludwigia linifolia</i> | | | | | | | |
| <i>Ludwigia mertima</i> | | | | | | | |
| <i>Ludwigia microcarpa</i> | | | | | | | |
| <i>Ludwigia octovalvis</i> | | | | | | | |
| <i>Lucknowia pilosa</i> | | | | | | | |
| <i>Ludwigia repens</i> | | | | | | | |
| <i>Ludwigia virgata</i> | | | | | | | |
| <i>Lycopodiella appressa</i> | Yes | Wet | 1-100 | 10,001-100,000 | South, Central | Asia | BW 218 |
| <i>Lycopodium microphyllum</i> | | | | | | | |
| <i>Lyonia fruticosa</i> | | | | | | | |
| <i>Lyonia lucida</i> | | | | | | | |
| <i>Lythrum alatum</i> var. <i>lanceolatum</i> | | | | | | | |
| <i>Marshallia tenuifolia</i> | | | | | | | |
| <i>Mecardonia acuminata</i> ssp. <i>peninsularis</i> | Yes | Wet | <1 | 101-1,000 | South, Central | Australia | BW 260 |
| <i>Melaleuca quinquenervia</i> | | | | | | | |
| <i>Melanthera nivea</i> | | | | | | | |
| <i>Mikania scandens</i> | | | | | | | |
| <i>Mitella petiolata</i> | | | | | | | |
| <i>Mitreola sessilifolia</i> | | | | | | | |
| <i>Muhlenbergia capillaris</i> | | | | | | | |
| <i>Musa x paradisiaca</i> | No | Mesic | <1 | 1-100 | South, Central, North | Asia | |
| <i>Myrica cerifera</i> | | | | | | | |
| <i>Nemastylis floridana</i> | | | | | | | |
| <i>Nephrolepis cordifolia</i> | Yes | Mesic | <1 | 101-1,000 | Throughout | Unknown | |
| <i>Nephrolepis exaltata</i> | | | | | | | |
| <i>Nuphar lutea</i> ssp. <i>advena</i> | | | | | | | |
| <i>Nymphaea odorata</i> | | | | | | | |
| <i>Nyctoides aquatica</i> | | | | | | | |

Pal-Mar - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|---------|----------------|---------------|------------------|---------------|
| <i>Osmunda cinnamomea</i> | | | | | | | |
| <i>Osmunda regalis</i> var. <i>spectabilis</i> | | | | | | | |
| <i>Oxypolis filiformis</i> | | | | | | | |
| <i>Panicum hemitomon</i> | | | | | | | |
| <i>Panicum maximum</i> | No | Mesic | <1 | 101-1,000 | Throughout | Africa | |
| <i>Panicum repens</i> | Yes | Wat | 1-100 | 10,001-100,000 | Throughout | Tropical America | |
| <i>Panicum rigidulum</i> | | | | | | | |
| <i>Panicum tenerum</i> | | | | | | | |
| <i>Panicum venosum</i> | | | | | | | |
| <i>Panicum virgatum</i> | | | | | | | |
| <i>Paspalidium geminatum</i> | | | | | | | |
| <i>Paspalum caespitosum</i> | | | | | | | |
| <i>Paspalum monostachyum</i> | | | | | | | |
| <i>Paspalum notatum</i> | No | Mesic | 1-100 | 101-1,000 | Throughout | South America | |
| <i>Paspalum praecox</i> | | | | | | | |
| <i>Paspalum setaceum</i> | | | | | | | |
| <i>Paspalum urvillei</i> | No | Mesic | <1 | 101-1,000 | Throughout | South America | BW 259 |
| <i>Passiflora suberosa</i> | | | | | | | |
| <i>Pectis glaucescens</i> | | | | | | | |
| <i>Pennisetum purpureum</i> | No | Mesic | <1 | 1-100 | Throughout | Africa | |
| <i>Pensea palustris</i> | | | | | | | |
| <i>Phalaris aureum</i> | | | | | | | |
| <i>Phragmites australis</i> | | | | | | | |
| <i>Phyla nodiflora</i> | | | | | | | |
| <i>Physalis walteri</i> | | | | | | | |
| <i>Pilobolus tigrinus</i> | | | | | | | |
| <i>Pinguicula</i> sp. | | | | | | | |
| <i>Pinus elliottii</i> var. <i>densa</i> | | | | | | | |
| <i>Piriqueta caroliniensis</i> | | | | | | | |
| <i>Plantanthera nivea</i> | | | | | | | |
| <i>Pluchea foetida</i> | | | | | | | |
| <i>Pluchea rosea</i> | | | | | | | |
| <i>Polygonia baldwinii</i> | | | | | | | |
| <i>Polygonia crucifera</i> | | | | | | | |
| <i>Polygonia grandiflora</i> | | | | | | | |
| <i>Polygonia lutea</i> | | | | | | | |
| <i>Polygonia ramosa</i> | | | | | | | |
| <i>Polygonia rugelii</i> | | | | | | | |
| <i>Polygonum hydropiperoides</i> | | | | | | | |
| <i>Polygonum punctatum</i> | | | | | | | |
| <i>Polygonum procumbens</i> | | | | | | | |
| <i>Pontederia cordata</i> | | | | | | | |
| <i>Proserpinaca palustris</i> | | | | | | | |
| <i>Proserpinaca pectinata</i> | | | | | | | |
| <i>Pteridium aquilinum</i> var. <i>caudatum</i> | | | | | | | |
| <i>Pteridium aquilinum</i> var. <i>pseudocaudatum</i> | | | | | | | |
| <i>Pterocaulon cynosbatioides</i> | | | | | | | |
| <i>Quercus minima</i> | | | | | | | |

Pal-Mar - Preliminary List of Vascular Plants

Bold = Exotic, *italicics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|----------------|-----------------------|---------------------|---------|
| <i>Rapanea punctata</i> | | | | | | | |
| <i>Rhexia mariana</i> | | | | | | | |
| <i>Rhus nuttallii</i> | | | | | | | |
| <i>Rhus copallinaum</i> | | | | | | | |
| <i>Rhynchospora repens</i> | No | Mesic | 1-100 | 10,001-100,000 | Throughout | South Africa | BW 250 |
| <i>Rhynchospora baldwinii</i> | | | | | | | |
| <i>Rhynchospora beriviseta</i> | | | | | | | |
| <i>Rhynchospora cephalantha</i> | | | | | | | |
| <i>Rhynchospora chapmanii</i> | | | | | | | |
| <i>Rhynchospora ciliolans</i> | | | | | | | |
| <i>Rhynchospora divergens</i> | | | | | | | |
| <i>Rhynchospora fascicularis</i> | | | | | | | |
| <i>Rhynchospora filifolia</i> | | | | | | | |
| <i>Rhynchospora hansei</i> | | | | | | | |
| <i>Rhynchospora inundata</i> | | | | | | | |
| <i>Rhynchospora laetifolia</i> | | | | | | | |
| <i>Rhynchospora microcarpa</i> | | | | | | | |
| <i>Rhynchospora microcephala</i> | | | | | | | |
| <i>Rhynchospora nitens</i> | | | | | | | |
| <i>Rhynchospora odorata</i> | | | | | | | |
| <i>Rhynchospora ratiflora</i> | | | | | | | |
| <i>Rhynchospora tracyi</i> | | | | | | | |
| <i>Sabal palmetto</i> | | | | | | | |
| <i>Sabatia bartramii</i> | | | | | | | |
| <i>Sabatia grandiflora</i> | | | | | | | |
| <i>Sabatia stellaris</i> | | | | | | | |
| <i>Saccharum giganteum</i> | | | | | | | |
| <i>Sacciolepis indica</i> | Yes | Mesic | 1-100 | >100,000 | Throughout | India | BW 258 |
| <i>Sacciolepis striata</i> | | | | | | | |
| <i>Sagittaria lancifolia</i> | | | | | | | |
| <i>Saxifrage caroliniana</i> | | | | | | | |
| <i>Samolus ebracteatus</i> | | | | | | | |
| <i>Sarcostemma clausum</i> | | | | | | | |
| <i>Schinus terebinthifolius</i> | Yes | Mesic | 1-100 | 10,001-100,000 | South, Central, North | Brazil and Paraguay | BW 248 |
| <i>Schizachyrium rhizomatum</i> | | | | | | | |
| <i>Schoenoliton albidiflorum</i> | | | | | | | |
| <i>Scolearius nigricans</i> | | | | | | | |
| <i>Scirpus hirtella</i> | | | | | | | |
| <i>Scirpus reticulans</i> | | | | | | | |
| <i>Scirpus triglomerata</i> | | | | | | | |
| <i>Scirpus verticillata</i> | | | | | | | |
| <i>Scoparia dulcis</i> | | | | | | | |
| <i>Senecio obtusifolia</i> | | | | | | | |
| <i>Sennaya repens</i> | | | | | | | |
| <i>Sebania herbacea</i> | | | | | | | |
| <i>Setaria parviflora</i> | | | | | | | |
| <i>Sida acuta</i> | | | | | | | |
| <i>Sida rhombifolia</i> | | | | | | | |
| <i>Sisyrinchium angustifolium</i> | | | | | | | |

Pai-Mar - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|--------------|---------------------------|------------------|---------|
| <i>Smilax auriculata</i> | | | | | | | |
| <i>Smilax laurifolia</i> | | | | | | | |
| <i>Solidago fistulosa</i> | | | | | | | |
| <i>Solidago gigantea</i> | | | | | | | |
| <i>Solidago stricta</i> | | | | | | | |
| <i>Sorghastrum secundatum</i> | | | | | | | |
| <i>Spermacoce assurgens</i> | | | | | | | |
| <i>Spermacoce verticillata</i> | Yes | Mesic | <1 | 101-1,000 | South, Central, North | Africa | |
| <i>Sporobolus indicus</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Tropical Asia | GB 1053 |
| <i>Sporobolus indicus</i> var. <i>pyramidalis</i> | No | Mesic | 1-100 | >100,000 | South, Central, Panhandle | Tropical America | BW 251 |
| <i>Sporobolus junceus</i> | | | | | | | |
| <i>Stenandrium dulce</i> | | | | | | | |
| <i>Stenotaphrum secundatum</i> | | | | | | | |
| <i>Stillingia aquatica</i> | | | | | | | |
| <i>Syngonanthus flavidulus</i> | | | | | | | |
| <i>Taxodium ascendens</i> | | | | | | | |
| <i>Taxodium distichum</i> | | | | | | | |
| <i>Thalia geniculata</i> | | | | | | | |
| <i>Thelypteris intermedia</i> | | | | | | | |
| <i>Thelypteris kunthii</i> | | | | | | | |
| <i>Tillandsia balbisiana</i> | | | | | | | |
| <i>Tillandsia fasciculata</i> var. <i>densispica</i> | | | | | | | |
| <i>Tillandsia flexuosa</i> | | | | | | | |
| <i>Tillandsia paucifolia</i> | | | | | | | |
| <i>Tillandsia recurvata</i> | | | | | | | |
| <i>Tillandsia setacea</i> | | | | | | | |
| <i>Tillandsia usneoides</i> | | | | | | | |
| <i>Tillandsia utriculata</i> | | | | | | | |
| <i>Toxicodendron radicans</i> | | | | | | | |
| <i>Tradescantia pallida</i> | No | Mesic | <1 | 1-100 | South, Central | Mexico | BW 238 |
| <i>Tripsacum dactyloides</i> | | | | | | | |
| <i>Typha domingensis</i> | | | | | | | |
| <i>Typha latifolia</i> | | | | | | | |
| <i>Urena lobata</i> | | | | | | | |
| <i>Urticaria comuta</i> | | | | | | | |
| <i>Urticaria fibrosa</i> | | | | | | | |
| <i>Urticaria foliosa</i> | | | | | | | |
| <i>Urticaria purpurea</i> | | | | | | | |
| <i>Urticaria resupinata</i> | | | | | | | |
| <i>Urticaria subulata</i> | | | | | | | |
| <i>Vaccinium myrsinifolium</i> | | | | | | | |
| <i>Verbena virginica</i> | | | | | | | |
| <i>Vernonia blodgettii</i> | | | | | | | |
| <i>Vigna luteola</i> | | | | | | | |
| <i>Vitis rotundifolia</i> | | | | | | | |
| <i>Vitis shuttleworthii</i> | | | | | | | |
| <i>Vitaria lineata</i> | | | | | | | |
| <i>Wedelia trilobata</i> | No | Mesic | <1 | 1-100 | Throughout | Tropical America | BW 224 |

Pal-Mar - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|----------------|---------------|------------------------|---------|
| Woodwardia virginica | | | | | | | |
| <i>Xyris ambigua</i> | | | | | | | |
| <i>Xyris difformis</i> , var. <i>floridana</i> | | | | | | | |
| <i>Xyris elliottii</i> | | | | | | | |
| <i>Xyris flabelliformis</i> | | | | | | | |
| <i>Xyris jupicai</i> | Yes | Wet | <1 | 10,001-100,000 | Throughout | Tropical America | BW 247 |
| <i>Xyris smalliana</i> | No | Mesic | <1 | 1-100 | Throughout | West Indies and Mexico | |
| <i>Yucca aloifolia</i> | | | | | | | |
| <i>Yucca filamentosa</i> | | | | | | | |

Pal-Mar - Endangered and Threatened Taxa

| | State Status | Habitat | Acreage | Pop. Size | # Stations | Status | Natural Range | Voucher |
|--|--------------|---------|-----------|--------------|------------|--------|--|------------------|
| <i>Eulophia alta</i> | T | Wet | <1 | 1-100 | Throughout | Stable | Florida, West Indies, Central and South America | BW 680 |
| <i>Lilium catesbaei</i> | T | Mesic | 1-100 | 101-1,000 | Throughout | Stable | Va. to La. and FL. | BW 698, 701, 744 |
| <i>Nemastylis floridana</i> | E | Wet | 101-1,000 | 101-1,000 | Throughout | Stable | South and Central Florida, north to Flagler County | |
| <i>Piatanthera nivea</i> | T | Wet | 1-100 | 1,001-10,000 | Throughout | Stable | Fla. to Tex., Ark., and N.J. | |
| <i>Tillandsia balbisiana</i> | T | Wet | 1-100 | 1,001-10,000 | Throughout | Stable | South and Central Florida, West Indies and Mexico to Colombia and Venezuela. | BW 231 |
| <i>Tillandsia fasciculata</i> var. <i>densispica</i> | E | Wet | 1-100 | 1,001-10,000 | Throughout | Stable | Florida to Central America | BW 233 |
| <i>Tillandsia flexuosa</i> | E | Wet | 1-100 | 101-1,000 | Throughout | Stable | South and Central Florida, West Indies, Panama, northern South America. | |
| <i>Tillandsia utriculata</i> | E | Wet | 1-100 | 1,001-10,000 | Throughout | Stable | Georgia and Florida, West Indies, Mexico, Northern South America, Venezuela. | |

Six Mile Cypress Preserve - Preliminary List of Vascular Plants

Bold = Exotic; *italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|---------|-----------|----------------|---------------|---------|
| | No | Mesic | <1 | 1-100 | South | Australia | GB 958 |
| Acacia auriculiformis | | | | | | | |
| Acalypha gracilens | | | | | | | |
| Acer rubrum | | | | | | | |
| Acemella oppositifolia var. <i>repens</i> | | | | | | | |
| Acrosichium danaeifolium | | | | | | | |
| Aeschynomene americana | | | | | | | |
| Agalinis fasciata | | | | | | | |
| Agalinis linifolia | | | | | | | |
| Aletris lutea | | | | | | | |
| Alternanthera philoxeroides | Yes | Wet | 1-100 | 101-1,000 | Throughout | South America | |
| Amaranthus australis | | | | | | | |
| Ambrosia artemisiifolia | | | | | | | |
| Ammannia latifolia | | | | | | | |
| Aniseiopsis arborea | | | | | | | |
| Amphicarpum muhlenbergianum | | | | | | | |
| Anachellium cochleatum var. <i>triandrum</i> | | | | | | | |
| Andropogon glomeratus var. <i>pumilus</i> | | | | | | | |
| Andropogon virginicus var. <i>decipiens</i> | | | | | | | |
| Andropogon virginicus var. <i>glaucus</i> | | | | | | | |
| Annona glabra | | | | | | | |
| Aristida beyrichiana | | | | | | | |
| Aristida palustris | | | | | | | |
| Aristida patula | | | | | | | |
| Aristida purpurascens | | | | | | | |
| Aristida spiciformis | | | | | | | |
| Asclepias incarnata | | | | | | | |
| Asclepias lanceolata | | | | | | | |
| Asimina reticulata | | | | | | | |
| Aster bracei | | | | | | | |
| Aster carolinianus | | | | | | | |
| Aster subulatus | | | | | | | |
| Axonopus fissifolius | | | | | | | |
| Axonopus furcatus | | | | | | | |
| Azolla caroliniana | | | | | | | |
| Baccharis angustifolia | | | | | | | |
| Baccharis glomeruliflora | | | | | | | |
| Baccharis halimifolia | | | | | | | |
| Bacopa caroliniana | | | | | | | |
| Bacopa monnieri | | | | | | | |
| Beffaria racemosa | | | | | | | |
| Bidens alba var. <i>radiata</i> | | | | | | | |
| Bigelowia nudata ssp. <i>australis</i> | Yes | Mesic | <1 | 1-100 | South, Central | Tropical Asia | |
| Bischofia javanica | | | | | | | |
| Blechnum semiaquatum | | | | | | | |

Six Mile Cypress Preserve - Preliminary List of Vascular Plants

Bold = Exotic; Italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|---------|-----------|----------------|---------------------------|---------|
| <i>Boehmeria cylindrica</i> | | | | | | | |
| <i>Bucinaria americana</i> | | | | | | | |
| <i>Burmannia capitata</i> | | | | | | | |
| <i>Calliandra americana</i> | | | | | | | |
| <i>Campyloneurum phyllidoides</i> | | | | | | | |
| <i>Canna flaccida</i> | | | | | | | |
| <i>Carex gigantea</i> | | | | | | | |
| <i>Carex lupulina</i> | | | | | | | |
| <i>Carex vexans</i> | | | | | | | |
| <i>Carphephorus conybosinus</i> | | | | | | | |
| <i>Carphephorus odoratissimus</i> | | | | | | | |
| <i>Carya aquatica</i> | | | | | | | |
| <i>Cassystha filiformis</i> | | | | | | | |
| <i>Cenchrus incertus</i> | | | | | | | |
| <i>Centella astatica</i> | | | | | | | |
| <i>Cephalanthus occidentalis</i> | | | | | | | |
| <i>Chamaecrista nictitans</i> var. <i>aspera</i> | | | | | | | |
| <i>Chamaesyce hirta</i> | | | | | | | |
| <i>Chamaesyce hypericifolia</i> | | | | | | | |
| <i>Chamaesyce hyssopifolia</i> | | | | | | | |
| <i>Chamaesyce ophthalmica</i> | | | | | | | |
| <i>Chaptalia tomentosa</i> | | | | | | | |
| <i>Chrysopogon pauciflorus</i> | | | | | | | |
| <i>Cirsium horridulum</i> | | | | | | | |
| <i>Cladodium jamaicense</i> | | | | | | | |
| <i>Coelorachis rugosa</i> | | | | | | | |
| <i>Commelinia diffusa</i> | | | | | | | |
| <i>Commelinia diffusa</i> var. <i>gigas</i> | | | | | | | |
| <i>Conodinum coelestinum</i> | | | | | | | |
| <i>Conyzia canadensis</i> var. <i>pusilla</i> | | | | | | | |
| <i>Cotopsis floridana</i> | | | | | | | |
| <i>Cotopsis leavenworthii</i> | | | | | | | |
| <i>Critmum americanum</i> | | | | | | | |
| <i>Crotalaria pallida</i> var. <i>obovata</i> | No | Mesic | <1 | 1-100 | Throughout | Africa | |
| <i>Crotalaria rotundifolia</i> | | | | | | | |
| <i>Croton glandulosus</i> var. <i>glandulosus</i> | | | | | South, Central | Australia | |
| <i>Cupaniopsis anacardioides</i> | Yes | Mesic | <1 | 1-100 | Throughout | South and Central America | |
| <i>Cuphea carthagenensis</i> | No | Wet | <1 | 101-1,000 | | | |
| <i>Cuscuta pentagona</i> | | | | | | | |
| <i>Cynanchum scoparium</i> | | | | | | | |
| <i>Cynodon dactylon</i> | | | | | | | |
| <i>Cyperus articulatus</i> | | | | | | | |
| <i>Cyperus croceus</i> | | | | | | | |
| <i>Cyperus distinctus</i> | | | | | | | |

BW 520

Six Mile Cypress Preserve - Preliminary List of Vascular Plants

Bold = Exotic; *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|--------------|---------------|-------------------|---------|
| <i>Cyperus erythrorhizos</i> | No | Mesic | <1 | 101-1,000 | Throughout | Old World | |
| <i>Cyperus esculentus</i> | No | Mesic | <1 | 101-1,000 | Throughout | Old World | BW 497 |
| <i>Cyperus haspan</i> | | | | | | | |
| <i>Cyperus ligularis</i> | | | | | | | |
| <i>Cyperus odoratus</i> | | | | | | | |
| <i>Cyperus polystachyos</i> | | | | | | | |
| <i>Cyperus retroflexus</i> | | | | | | | |
| <i>Cyperus surinamensis</i> | | | | | | | |
| <i>Dactyloctenium aegyptium</i> | No | Mesic | <1 | 101-1,000 | Throughout | Old World Tropics | |
| <i>Desmodium incanum</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Old World | |
| <i>Desmodium triflorum</i> | No | Mesic | <1 | 1,001-10,000 | Throughout | Old World | |
| <i>Dichanthelium aciculare</i> | | | | | | | |
| <i>Dichanthelium commutatum</i> | | | | | | | |
| <i>Dichanthelium dichotomum</i> | | | | | | | |
| <i>Dichanthelium erectifolium</i> | | | | | | | |
| <i>Dichanthelium leucotrich</i> | | | | | | | |
| <i>Dichanthelium portoricense</i> | | | | | | | |
| <i>Dichanthelium strigosum</i> var. <i>glabrescens</i> | | | | | | | |
| <i>Digitaria ciliaris</i> | | | | | | | |
| <i>Digitaria villosa</i> | | | | | | | |
| <i>Dioclea virginiana</i> | | | | | | | |
| <i>Drosera capillaris</i> | | | | | | | |
| <i>Dyschoriste oblongifolia</i> | | | | | | | |
| <i>Echinochloa paludigena</i> | | | | | | | |
| <i>Echinochloa walteri</i> | | | | | | | |
| <i>Eclipta prostrata</i> | Yes | Wet | 1-10 | 1,001-10,000 | Throughout | Brazil | |
| <i>Eichornia crassipes</i> | Yes | Wet | 1-10 | 1,001-10,000 | Throughout | Brazil | BW 528 |
| <i>Eleocharis baldwinii</i> | | | | | | | |
| <i>Eleocharis cellososa</i> | | | | | | | |
| <i>Eleocharis geniculata</i> | | | | | | | |
| <i>Eleocharis nigrescens</i> | Yes | Mesic | 1-10 | 1,001-10,000 | Throughout | Pantropical | |
| <i>Elyonurus tripsacoides</i> | | | | | | | |
| <i>Elytraria carolinensis</i> var. <i>angustifolia</i> | | | | | | | |
| <i>Emilia fosbergii</i> | No | Mesic | <1 | 101-1,000 | Throughout | Old World | BW 519 |
| <i>Encyclia tampensis</i> | | | | | | | |
| <i>Epidendrum rigidum</i> | | | | | | | |
| <i>Eragrostis ciliaris</i> | No | Mesic | <1 | 1-100 | Throughout | New World Tropics | |
| <i>Eragrostis elatior</i> | | | | | | | |
| <i>Eragrostis virginica</i> | | | | | | | |
| <i>Erechtites hieracifolia</i> | | | | | | | |
| <i>Erigeron quercifolius</i> | | | | | | | |
| <i>Erigeron vernus</i> | | | | | | | |
| <i>Eriogonum decangulare</i> | | | | | | | |

Six Mile Cypress Preserve - Preliminary List of Vascular Plants

Bold = Exotic; *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---------------------------------|------------|---------|---------|--------------|----------------|------------------|---------|
| <i>Eriochloa michauxii</i> | | | | | | | |
| <i>Eryngium baldwinii</i> | | | | | | | |
| <i>Eryngium yuccifolium</i> | | | | | | | |
| <i>Euphorbia alta</i> | | | | | | | |
| <i>Eupatorium capillifolium</i> | | | | | | | |
| <i>Eupatorium leptophyllum</i> | | | | | | | |
| <i>Eupatorium milianoides</i> | | | | | | | |
| <i>Eupatorium mohrii</i> | | | | | | | |
| <i>Eupatorium rotundifolium</i> | | | | | | | |
| <i>Euphorbia polyphylla</i> | | | | | | | |
| <i>Eustachys glauca</i> | | | | | | | |
| <i>Eustachys petraea</i> | | | | | | | |
| <i>Euthamia caroliniana</i> | | | | | | | |
| <i>Evolvulus sericeus</i> | | | | | | | |
| <i>Ficus aurea</i> | Yes | Mesic | <1 | 1-100 | South, Central | Old World | BW 179 |
| <i>Ficus microcarpa</i> | | | | | | | |
| <i>Fimbristylis autumnalis</i> | | | | | | | |
| <i>Fimbristylis puberula</i> | | | | | | | |
| <i>Fimbristylis spadicea</i> | | | | | | | |
| <i>Flaveria linearis</i> | | | | | | | |
| <i>Fraxinus caroliniana</i> | | | | | | | |
| <i>Fultrena brevirostra</i> | | | | | | | |
| <i>Fultrena pumila</i> | | | | | | | |
| <i>Fultrena scirpoidea</i> | | | | | | | |
| <i>Galectia elliottii</i> | | | | | | | |
| <i>Galactia regularis</i> | | | | | | | |
| <i>Gaura angustifolia</i> | | | | | | | |
| <i>Gaultheria dumosa</i> | No | Mesic | <1 | 1-100 | Throughout | Tropical America | BW 525 |
| <i>Gomphrena serrata</i> | | | | | | | |
| <i>Gratiola ramosa</i> | | | | | | | |
| <i>Habenaria quinqueseta</i> | | | | | | | |
| <i>Harrisia filiformis</i> | | | | | | | |
| <i>Hedychium procumbens</i> | | | | | | | |
| <i>Hedychium uniflora</i> | | | | | | | |
| <i>Helianthus debilis</i> | | | | | | | |
| <i>Heleocephalum polypodium</i> | No | Mesic | <1 | 1-100 | South, Central | Pantropical | BW 172 |
| <i>Hemarthria altissima</i> | | | | | | | |
| <i>Hererotheca subaxillaris</i> | | | | | | | |
| <i>Hieracium megacephalon</i> | | | | | | | |
| <i>Hydrocotyle umbellata</i> | | | | | | | |
| <i>Hydrocotyle verticillata</i> | | | | | | | |
| <i>Hydroclea corymbosa</i> | Yes | Wet | 1-100 | 1,001-10,000 | South, Central | West Indies | GB 964 |
| <i>Hypéricum brachyphyllum</i> | | | | | | | |

Six Mile Cypress Preserve - Preliminary List of Vascular Plants

Bold = Exotic; *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|-----------|-----------------------|---------------|---------|
| <i>Hypericum cistifolium</i> | | | | | | | |
| <i>Hypericum fasciculatum</i> | | | | | | | |
| <i>Hypericum hypericoides</i> | | | | | | | |
| <i>Hypericum reducillum</i> | | | | | | | |
| <i>Hypericum tetrapetalum</i> | | | | | | | |
| <i>Hypoxis curtissii</i> | | | | | | | |
| <i>Hypoxis luncea</i> | | | | | | | |
| <i>Hypolis alata</i> | | | | | | | |
| <i>Ilex cassine</i> | | | | | | | |
| <i>Ilex glabra</i> | | | | | | | |
| <i>Ipomoea alba</i> | | | | | | | |
| <i>Ipomoea sagittata</i> | | | | | | | |
| <i>Iris hexagona</i> | | | | | | | |
| <i>Iva microcephala</i> | | | | | | | |
| <i>Juncus effusus</i> var. <i>solutus</i> | | | | | | | |
| <i>Juncus marginatus</i> | | | | | | | |
| <i>Juncus megacephalus</i> | | | | | | | |
| <i>Juncus polycerathus</i> | | | | | | | |
| <i>Juncus scirpoides</i> | | | | | | | |
| <i>Juniperus virginiana</i> | | | | | | | |
| <i>Justicia angusta</i> | | | | | | | |
| <i>Lachnocaulon anceps</i> | - | | | | | | |
| <i>Lachnocaulon engleri</i> | | | | | | | |
| <i>Lechea torreyi</i> | | | | | | | |
| <i>Leersia hexandra</i> | | | | | | | |
| <i>Lemna valdiviana</i> | | | | | | | |
| <i>Leptochloa fascicularis</i> | No | Mesic | <1 | 1-100 | South, Central, North | Caribbean? | |
| <i>Leucena leucocephala</i> | | | | | | | |
| <i>Liatris gracilis</i> | | | | | | | |
| <i>Liatris tenuifolia</i> | | | | | | | |
| <i>Lilium catesbeianum</i> | No | Mesic | <1 | 101-1,000 | Throughout | Indo-Malaya | BW 764 |
| <i>Lindernia crustacea</i> | | | | | | | |
| <i>Lindernia dubia</i> var. <i>anagallidea</i> | | | | | | | |
| <i>Linum medium</i> var. <i>texanum</i> | | | | | | | |
| <i>Lipocarpha micrantha</i> | | | | | | | |
| <i>Liquidambar styraciflua</i> | | | | | | | |
| <i>Lobelia glandulosa</i> | | | | | | | |
| <i>Lobelia paludosa</i> | | | | | | | |
| <i>Ludwigia alata</i> | | | | | | | |
| <i>Ludwigia curtissii</i> | | | | | | | |
| <i>Ludwigia maritima</i> | | | | | | | |
| <i>Ludwigia microcarpa</i> | | | | | | | |
| <i>Ludwigia octovalvis</i> | | | | | | | |
| <i>Ludwigia peruviana</i> | | | | | | | |

Six Mile Cypress Preserve - Preliminary List of Vascular Plants

Bold = Exotic; Italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|----------------|----------------|------------------|---------------|
| <i>Ludwigia repens</i> | | | | | | | |
| <i>Ludwigia simpsonii</i> | | | | | | | |
| <i>Lycopus rubellus</i> | | | | | | | |
| <i>Lygodesmia achylla</i> | | | | | | | |
| <i>Lygodium microphyllum</i> | Yes | Wet | <1 | 1-100 | South, Central | Asia | BW 510 |
| <i>Lyonia fruticosa</i> | | | | | | | |
| <i>Lythrum alatum</i> var. <i>lanceolatum</i> | | | | | | | |
| <i>Magnolia virginiana</i> | | | | | | | |
| <i>Maiaxis spicata</i> | | | | | | | |
| <i>Mecardonia acuminata</i> ssp. <i>peninsularis</i> | | | | | | | |
| <i>Melaleuca quinquenervia</i> | Yes | Wet | 1-100 | 10,001-100,000 | South, Central | Australia | |
| <i>Melanthera nivea</i> | | | | | | | |
| <i>Melochia spicata</i> | | | | | | | |
| <i>Melothria pendula</i> | | | | | | | |
| <i>Micranthemum umbrosum</i> | | | | | | | |
| <i>Mikania scandens</i> | | | | | | | |
| <i>Mitreola petiolata</i> | | | | | | | |
| <i>Mitreola sessiliifolia</i> | | | | | | | |
| <i>Muhlenbergia capillaris</i> | No | Mesic | <1 | 1-100 | Throughout | Asia | BW 774 |
| <i>Murdannia nudiflora</i> | | | | | | | |
| <i>Myrica cerifera</i> | | | | | | | |
| <i>Myriophyllum heterophyllum</i> | | | | | | | |
| <i>Nephrolepis cordifolia</i> | Yes | Mesic | <1 | 101-1,000 | Throughout | Unknown | |
| <i>Nephrolepis exaltata</i> | | | | | | | |
| <i>Nothoscordum bivalve</i> | | | | | | | |
| <i>Nymphaea elegans</i> | | | | | | | |
| <i>Nyssa sylvatica</i> var. <i>biflora</i> | | | | | | | |
| <i>Osmunda regalis</i> var. <i>spectabilis</i> | | | | | | | |
| <i>Oxalis corniculata</i> | | | | | | | |
| <i>Oxybaphus filiformis</i> | | | | | | | |
| <i>Panicum anceps</i> | | | | | | | |
| <i>Panicum dichotomiflorum</i> | | | | | | | |
| <i>Panicum hemitomon</i> | | | | | | | |
| <i>Panicum hians</i> | | | | | | | |
| <i>Panicum repens</i> | Yes | Wet | 1-100 | 1,001-10,000 | Throughout | Tropical America | BW 524 |
| <i>Panicum rigidulum</i> | | | | | | | |
| <i>Panicum tenerum</i> | | | | | | | |
| <i>Parthenocissus quinquefolia</i> | | | | | | | |
| <i>Paspalum laeve</i> | | | | | | | |
| <i>Paspalum monostachyum</i> | No | Mesic | 1-100 | 1,001-10,000 | Throughout | South America | |
| <i>Paspalum notatum</i> | | | | | | | |
| <i>Paspalum repens</i> | | | | | | | |
| <i>Paspalum setaceum</i> | No | Mesic | <1 | 101-1,000 | Throughout | South America | |
| <i>Paspalum urvillei</i> | No | Mesic | | | | | BW 527 |

Six Mile Cypress Preserve - Preliminary List of Vascular Plants

Bold = Exotic; *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|-----------|---------------|----------------|------------------|
| <i>Passiflora suberosa</i> | | | | | | | |
| <i>Pectis glaucescens</i> | | | | | | | |
| <i>Pectis linearifolia</i> | | | | | | | |
| <i>Persea palustris</i> | | | | | | | |
| <i>Phlebodium aureum</i> | | | | | | | |
| <i>Phoradendron leucarpum</i> | | | | | | | |
| <i>Phyla nodiflora</i> | | | | | | | |
| <i>Phyllanthus caroliniensis</i> var. <i>saxicola</i> | No | | Mesic | <1 | 1-100 | Throughout | |
| <i>Phyllanthus urinaria</i> | | | | | | | |
| <i>Physalis walteri</i> | | | | | | | |
| <i>Physostegia purpurea</i> | | | | | | | |
| <i>Pileolepis rigida</i> | | | | | | | |
| <i>Pinguicula pumila</i> | | | | | | | |
| <i>Pinguicula</i> sp. | | | | | | | |
| <i>Pinus elliottii</i> var. <i>densa</i> | | | | | | | |
| <i>Piniella caroliniana</i> | | | | | | | |
| <i>Pistia stratiotes</i> | Yes | | Wet | 1-100 | >100,000 | Throughout | |
| <i>Pitchea odorata</i> | | | | | | | |
| <i>Plucephala rosea</i> | | | | | | | |
| <i>Polygonia baldwinii</i> | | | | | | | |
| <i>Polygonia grandiflora</i> | | | | | | | |
| <i>Polygonia incarnata</i> | | | | | | | |
| <i>Polygonia lutea</i> | | | | | | | |
| <i>Polygonia setacea</i> | | | | | | | |
| <i>Polygonella polygama</i> var. <i>polygama</i> | | | | | | | |
| <i>Polygonum densiflorum</i> | | | | | | | |
| <i>Polygonum hydrotriperoides</i> | | | | | | | |
| <i>Polygonum punctatum</i> | | | | | | | |
| <i>Polypodium polypodioides</i> var. <i>mitchauxiana</i> | | | | | | | |
| <i>Polypteron procumbens</i> | | | | | | | |
| <i>Polystachya concreta</i> | | | | | | | |
| <i>Panoleleria cordata</i> | | | | | | | |
| <i>Portulaca amilis</i> | No | | Mesic | <1 | 1-100 | Throughout | |
| <i>Proserpinaca palustris</i> | | | | | | | |
| <i>Proserpinaca pectinata</i> | | | | | | | |
| <i>Psiadium cattleyanum</i> | Yes | | Mesic | <1 | 1-100 | South, Central | Tropical America |
| <i>Psiotom nudum</i> | | | | | | | |
| <i>Psychotria sulzneri</i> | | | | | | | |
| <i>Pleidium aquilinum</i> var. <i>caudatum</i> | | | | | | | |
| <i>Pleidium aquilinum</i> var. <i>pseudocaudatum</i> | Yes | | Mesic | <1 | 101-1,000 | Throughout | Eastern Asia |
| <i>Pteris vittata</i> | | | | | | | |
| <i>Pterocaulon pycnostachyum</i> | | | | | | | |
| <i>Ptilimnium capillaceum</i> | | | | | | | |
| <i>Quercus laurifolia</i> | | | | | | | |

Six Mile Cypress Preserve - Preliminary List of Vascular Plants

Bold = Exotic; *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|-----------|--------------|-----------------------|---------------------|---------|
| <i>Quercus minima</i> | | | | | | | |
| <i>Quercus virginiana</i> | | | | | | | |
| <i>Rapanea punctata</i> | | | | | | | |
| <i>Rhexia mariana</i> | | | | | | | |
| <i>Rhexia nuttallii</i> | | | | | | | |
| <i>Rhynchospora repens</i> | No | Mesic | 1-100 | 101-1,000 | Throughout | South Africa | BW 524 |
| <i>Rhynchospora colorata</i> | | | | | | | |
| <i>Rhynchospora corniculata</i> | | | | | | | |
| <i>Rhynchospora divergens</i> | | | | | | | |
| <i>Rhynchospora fascicularis</i> | | | | | | | |
| <i>Rhynchospora fermaidii</i> | | | | | | | |
| <i>Rhynchospora intermedia</i> | | | | | | | |
| <i>Rhynchospora inundata</i> | | | | | | | |
| <i>Rhynchospora microcarpa</i> | | | | | | | |
| <i>Rhynchospora milacea</i> | | | | | | | |
| <i>Rhynchospora nitens</i> | | | | | | | |
| <i>Rhynchospora odorata</i> | | | | | | | |
| <i>Rhynchospora tracyi</i> | | | | | | | |
| <i>Rotala ramosior</i> | | | | | | | |
| <i>Rudbeckia hirta</i> | | | | | | | |
| <i>Ruellia</i> sp. | | | | | | | |
| <i>Rumex verticillatus</i> | | | | | | | |
| <i>Sabal palmetto</i> | | | | | | | |
| <i>Sabatia bartramii</i> | | | | | | | |
| <i>Sabatia brevifolia</i> | | | | | | | |
| <i>Sabatia calycina</i> | | | | | | | |
| <i>Sabatia stellaris</i> | | | | | | | |
| <i>Saccharum giganteum</i> | | | | | | | |
| <i>Sacciolepis indica</i> | Yes | Mesic | 1-100 | 1,001-10,000 | Throughout | India | BW 763 |
| <i>Sacciolepis striata</i> | | | | | | | |
| <i>Saccolia lanceolata</i> | | | | | | | |
| <i>Sagittaria filiformis</i> | | | | | | | |
| <i>Sagittaria graminea</i> var. <i>chapmanii</i> | | | | | | | |
| <i>Sagittaria lancifolia</i> | | | | | | | |
| <i>Salix caroliniana</i> | | | | | | | |
| <i>Savinnia minima</i> | | | | | | | |
| <i>Samolus ebracteatus</i> | | | | | | | |
| <i>Sarcostemma clausum</i> | | | | | | | |
| <i>Schiminthus terebinthifolius</i> | Yes | Mesic | 101-1,000 | >100,000 | South, Central, North | Brazil and Paraguay | BW 783 |
| <i>Schizachyrium thizomatum</i> | | | | | | | |
| <i>Schoenolirion albiflorum</i> | | | | | | | |
| <i>Schoenus nigricans</i> | | | | | | | |
| <i>Scirpus pungens</i> | | | | | | | |
| <i>Scirpus tabernaemontani</i> | | | | | | | |

Six Mile Cypress Preserve - Preliminary List of Vascular Plants

Bold = Exotic; *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|---------|----------------|-----------------------|----------------------|---------------|
| <i>Sciria baldwinii</i> | | | | | | | |
| <i>Sciria reticularis</i> | | | | | | | |
| <i>Sciria verticillata</i> | | | | | | | |
| <i>Scoparia dulcis</i> | | | | | | | |
| <i>Senna obtusifolia</i> | | | | | | | |
| <i>Senna pendula</i> var. <i>glabrata</i> | Yes | Wet | 1-100 | 1,001-10,000 | South, Central | South America | |
| <i>Serenoa repens</i> | | | | | | | |
| <i>Sesbania herbacea</i> | | | | | | | |
| <i>Selaria magna</i> | | | | | | | |
| <i>Selaria parviflora</i> | | | | | | | |
| <i>Sida acuta</i> | | | | | | | |
| <i>Sida cordifolia</i> | No | Mesic | <1 | 101-1,000 | South, Central, North | Tropical America | |
| <i>Sida elliptii</i> | | | | | | | |
| <i>Sida rhombifolia</i> | | | | | | | |
| <i>Sideroxylon redinatum</i> | | | | | | | |
| <i>Smilax auriculata</i> | | | | | | | |
| <i>Smilax bona-nox</i> | | | | | | | |
| <i>Smilax laurifolia</i> | | | | | | | |
| <i>Solanum americanum</i> | | | | | | | |
| <i>Solanum tampicense</i> | Yes | Wet | 1-100 | 10,001-100,000 | South, Central | Mexico | |
| <i>Solanum viarum</i> | No | Mesic | <1 | 1-100 | Throughout | Argentina and Brazil | |
| <i>Solidago fistulosa</i> | | | | | | | |
| <i>Solidago gigantea</i> | | | | | | | |
| <i>Solidago sempervirens</i> | | | | | | | |
| <i>Solidago stricta</i> | | | | | | | |
| <i>Sorghastrum secundum</i> | | | | | | | |
| <i>Spartina bakeri</i> | | | | | | | |
| <i>Spermacoce assurgens</i> | | | | | | | |
| <i>Spermacoce verticillata</i> | No | Mesic | <1 | 101-1,000 | South, Central, North | Africa | BW 521 |
| <i>Spiranthes odorata</i> | | | | | | | |
| <i>Spirodela polyrhiza</i> | | | | | | | |
| <i>Spirodela punctata</i> | Yes | Wet | 1-100 | >100,000 | Throughout | Old World | |
| <i>Sporobolus indicus</i> | No | Mesic | <1 | 101-1,000 | Throughout | Tropical Asia | |
| <i>Sporobolus junceus</i> | | | | | | | |
| <i>Stenandrium dulce</i> | | | | | | | |
| <i>Stillingia aquatica</i> | | | | | | | |
| <i>Stillingia sylvatica</i> | | | | | | | |
| <i>Sweetenia mahagoni</i> | | | | | | | |
| <i>Syngonanthus flavidulus</i> | | | | | | | |
| <i>Taxodium ascendens</i> | | | | | | | |
| <i>Taxodium distichum</i> | | | | | | | |
| <i>Tephrosia rugellii</i> | | | | | | | |
| <i>Teucrium canadense</i> | | | | | | | |
| <i>Thalia geniculata</i> | | | | | | | |

Six Mile Cypress Preserve - Endangered and Threatened Taxa

| State Status | Habitat | Acreage | Pop. Size | # Station | Status | Natural Range | Voucher |
|--------------|---------|-----------|--------------|-----------|-------------|---|---------|
| E | Wet | <1 | 1-100 | 1 | Decreasing? | Florida to South America | BW 175 |
| E | Wet | <1 | 101-1,000 | Scattered | Stable | Florida to South America | BW 531 |
| E | Wet | <1 | 1-100 | 1 | Decreasing? | Florida to South America | BW 176 |
| T | Wet | <1 | 1-100 | Scattered | Stable | Florida, West Indies, Central and South America | |
| T | Wet | <1 | 101-1,000 | 3 | Stable | Florida to Central America | BW 163 |
| T | Mesic | <1 | 1-100 | 2 | Stable | Va. to La. and F.L. | BW 749 |
| E | Wet | <1 | 1-100 | 1 | Decreasing? | Florida to South America | BW 177 |
| T | Wet | 101-1,000 | 1,001-10,000 | Scattered | Stable | Florida to South America | BW 507 |
| E | Wet | 101-1,000 | 1,001-10,000 | Scattered | Stable | Florida to Central America | |
| E | Wet | 101-1,000 | 1,001-10,000 | Scattered | Stable | Georgia to South America | |
| T | Wet | 1-100 | 1,001-10,000 | Scattered | Stable | Florida to South America | |

Water Conservation Area 2 - Preliminary List of Vascular Plants

Bold = Exotic, Italics = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|---------|--------------|---------------------------|------------------|--------------------|
| Acacia auriculiformis | No | Mesic | <1 | 101-1,000 | South | Australia | BW 154 |
| Acacia farnesiana | | | | | | | |
| <i>Acer rubrum</i> | | | | | | | |
| Acrostichum danaeifolium | | | | | | | |
| Amaranthus australis | | | | | | | |
| Ambrosia artemisiifolia | | | | | | | |
| Ammannia latifolia | | | | | | | |
| Ampelopsis arborea | | | | | | | |
| Andropogon glomeratus var. pumilus | | | | | | | |
| Annona glabra | No | Mesic | 1-100 | 1-100 | Throughout | Mexico | BW 306 |
| Antigonon leptopus | | | | | | | |
| Apocynum cannabinum | | | | | | | |
| Ardisia escallonioides | | | | | | | |
| Aristida purpurascens | | | | | | | |
| Aster carolinianus | | | | | | | |
| Aster dumosus | | | | | | | |
| Aster subulatus | | | | | | | |
| Baccharis angustifolia | | | | | | | |
| Bacopa monnieri | | | | | | | |
| Bidens alba var. radiata | | | | | | | |
| Bischoffia javanica | No | Mesic | <1 | 1-100 | South | Tropical Asia | BW 155, 297 |
| Blechnum serrulatum | | | | | | | |
| Boehmeria cylindrica | | | | | | | |
| Buchnera americana | | | | | | | |
| Bucida buceras | No | Mesic | <1 | 1-100 | South | Caribbean | BW 151 |
| Bursera simaruba | | | | | | | |
| Canna flaccida | | | | | | | |
| Cardiospermum halicacabum | No | Mesic | <1 | 1-100 | South, Central | American tropics | |
| Carica papaya | Yes | Mesic | 1-100 | 101-1,000 | South, Central | Tropical America | BW 296 |
| Caryota mitis | No | Mesic | <1 | 1-100 | South | Old World | |
| Casuarina equisetifolia | No | Mesic | 1-100 | 1,001-10,000 | South, Central, Panhandle | Australia | |
| Catharanthus roseus | No | Mesic | 1-100 | 1,001-10,000 | South, Central, Panhandle | Madagascar | |
| Celtis laevigata | | | | | | | |
| Cenchrus echinatus | | | | | | | |
| Cenchrus integer | | | | | | | |
| Centella asiatica | | | | | | | |
| Cephaelanthus occidentalis | | | | | | | |
| Ceratophyllum demersum | | | | | | | |
| Cestrum diurnum | No | Mesic | 1-100 | 1-100 | South, Central | West Indies | BW 153 |
| <i>Chamaesyce blodgettii</i> | | | | | | | |
| <i>Chamaesyce hirta</i> | | | | | | | |
| <i>Chamaesyce hypericifolia</i> | | | | | | | |
| <i>Chamaesyce hyssopifolia</i> | | | | | | | |

Water Conservation Area 2 - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|----------------------------------|------------|---------|--------------|----------------|-----------------------|--|---------|
| Emilia fosbergii | No | Mesic | <1 | 1,001-10,000 | Throughout | Old World | |
| Eragrostis atrovirens | Yes | Mesic | <1 | 10,001-100,000 | Throughout | Southeast Asia | |
| Eragrostis ciliaris | No | Mesic | <1 | 1,001-10,000 | Throughout | Coastal Plain, Fl. to Tex. and N.J. - (W.I., Mex., C.A., S.A.) | BW 149 |
| Eragrostis ellottii | | | | | | | |
| Erechtites hieracifolia | | | | | | | |
| Eremochloa ophiuroides | No | Mesic | 1-100 | >100,000 | Throughout | Southeast Asia | |
| Eriogonum quercifolius | | | | | | | |
| Eugenia uniflora | No | Mesic | <1 | 1-100 | South, Central | South America | |
| Eupatorium capillifolium | | | | | | | |
| Eupatorium serotinum | | | | | | | |
| Eustachys petraea | | | | | | | |
| Evolvulus sericeus | | | | | | | |
| Ficus aurea | | | | | | | |
| Ficus citrifolia | | | | | | | |
| Ficus microcarpa | Yes | Mesic | <1 | 1-100 | South | Old World | |
| Fimbristylis cymosa | | | | | | | |
| Flaveria linearis | | | | | | | |
| Fuliginea breviseta | | | | | | | |
| Gaura angustifolia | | | | | | | |
| Heliotropium angiospermum | | | | | | | |
| Heliotropium polypyllum | | | | | | | |
| Heterotheca subaxillaris | | | | | | | |
| Hibiscus grandiflorus | | | | | | | |
| Hydrilla verticillata | Yes | Wet | 1,001-10,000 | >100,000 | Throughout | Old World | |
| Hydrocotyle verticillata | | | | | | | |
| Hypenticum brachyphyllum | | | | | | | |
| Hypizia verticillata | No | Mesic | <1 | 1-100 | South, Central | Tropical America | |
| Ilex cassine | | | | | | | |
| Indigofera suffruticosa | No | Mesic | <1 | 101-1,000 | South, Central, North | New World Tropics | |
| Ipomoea alba | | | | | | | |
| Ipomoea sagittata | | | | | | | |
| Iresine diffusa | | | | | | | |
| Iva microcephala | | | | | | | |
| Juncus megacephalus | | | | | | | |
| Justicia angusta | | | | | | | |
| Kosteletzkyia virginica | | | | | | | |
| Lantana camara | No | Mesic | 1-100 | 1,001-10,000 | Throughout | Tropical America | |
| Lantana involucrata | | | | | | | |
| Leersia hexandra | | | | | | | |
| Lenna validiana | | | | | | | |
| Lepidium virginicum | | | | | | | |
| Lepiochloa fasciculata | No | Mesic | <1 | 1-100 | South, Central, North | Possibly Caribbean | |
| Lettuce leucocephala | | | | | | | |

Water Conservation Area 2 - Preliminary List of Vascular Plants

Bold = Exotic, *italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|---|------------|---------|--------------|--------------|---------------------------|---------------------------|-------------|
| <i>Ludwigia alata</i> | | | | | | | |
| <i>Ludwigia octovalvis</i> | | | | | | | |
| <i>Ludwigia peruviana</i> | | | | | | | |
| <i>Ludwigia repens</i> | | | | | | | |
| <i>Luziola fluitans</i> | | | | | | | |
| <i>Lysimoma sabiceu</i> | No | Mesic | <1 | 1-100 | South | Bahamas, Cuba, Hispaniola | |
| <i>Lythrum alatum</i> var. <i>lanceolatum</i> | No | Mesic | 1-100 | 1,001-10,000 | South, Central, Panhandle | Tropical America | BW 308 |
| <i>Macropilium lathyroides</i> | No | Mesic | | | | | |
| <i>Magnolia virginiana</i> | | | | | | | |
| <i>Melaleuca quinquenervia</i> | Yes | Wet | 1,001-10,000 | >100,000 | South, Central | Australia | |
| <i>Melothria pendula</i> | No | Mesic | <1 | 101-1,000 | Throughout | Tropical America | |
| <i>Merremia dissecta</i> | No | Mesic | | | | | |
| <i>Mikania scandens</i> | | | | | | | |
| <i>Mitreola petiolata</i> | | | | | | | |
| <i>Momordica charantia</i> | No | Mesic | <1 | 1,001-10,000 | South, Central, North | Old World Tropics | |
| <i>Morus rubra</i> | | | | | | | |
| <i>Muhlenbergia capillaris</i> | | | | | | | |
| <i>Musa x paradisiaca</i> | No | Mesic | <1 | 1-100 | South, Central | Asia | |
| <i>Myrica cerifera</i> | | | | | | | |
| <i>Najas guadalupensis</i> | | | | | | | |
| <i>Neuroleia reynaudiana</i> | No | Mesic | <1 | 1-100 | South | Southern Asia | BW 158 |
| <i>Nuphar lutea</i> ssp. <i>advena</i> | | | | | | | |
| <i>Nymphaea odorata</i> | | | | | | | |
| <i>Opismenus hirtellus</i> | | | | | | | |
| <i>Panicum dichotomiflorum</i> | | | | | | | |
| <i>Panicum dichotomiflorum</i> var. <i>bartowense</i> | | | | | | | |
| <i>Panicum hemitomon</i> | No | Mesic | 101-1,000 | 1,001-10,000 | Throughout | Africa | |
| <i>Panicum maximum</i> | Yes | Wet | 101-1,000 | >100,000 | Throughout | Tropical America | BW 146, 286 |
| <i>Panicum rigidulum</i> | | | | | | | |
| <i>Parietaria floridana</i> | | | | | | | |
| <i>Parthenium hysterophorus</i> | No | Mesic | 1-100 | 1,001-10,000 | Throughout | Tropical America | |
| <i>Pathenocissus quinquefolia</i> | | | | | | | |
| <i>Paspalidium geminalum</i> | | | | | | | |
| <i>Paspalum blodgettii</i> | | | | | | | |
| <i>Paspalum caespitosum</i> | | | | | | | |
| <i>Paspalum conjugatum</i> | | | | | | | |
| <i>Paspalum distichum</i> | | | | | | | |
| <i>Paspalum setaceum</i> | | | | | | | |
| <i>Paspalum urvillei</i> | No | Mesic | <1 | 101-1,000 | Throughout | South America | |
| <i>Passiflora suberosa</i> | | | | | | | |
| <i>Pectis glaucescens</i> | | | | | | | |
| <i>Pectis prostrata</i> | | | | | | | |

Water Conservation Area 2 - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|-----------|----------------|---------------------------|---|---------------|
| <i>Salix caroliniana</i> | | | | | | | |
| <i>Salvia occidentalis</i> | | | | | | | |
| <i>Salvia minima</i> | | | | | | | |
| <i>Sambucus canadensis</i> | | | | | | | |
| <i>Sarcostemma clausum</i> | | | | | | | |
| <i>Scirurus cemuis</i> | | | | | | | |
| <i>Schefflera actinophylla</i> | No | Mesic | <1 | 101-1,000 | South | Australia | |
| <i>Schinus terebinthifolius</i> | Yes | Mesic | 101-1,000 | >100,000 | South, Central, North | Brazil and Paraguay | BW 152 |
| <i>Schizachyrium gracile</i> | | | | | | | |
| <i>Schizachyrium sanguineum</i> var. <i>sanguineum</i> | | | | | | | |
| <i>Schoenus nigricans</i> | | | | | | | |
| <i>Scirpus labernaemontani</i> | | | | | | | |
| <i>Scoparia dulcis</i> | | | | | | | |
| <i>Senna ligustrina</i> | | | | | | | |
| <i>Senna obtusifolia</i> | No | Mesic | <1 | 101-1,000 | South, Central | South America | |
| <i>Senna pendula</i> var. <i>glabratia</i> | | | | | | | |
| <i>Sesbania herbacea</i> | | | | | | | |
| <i>Setaria magna</i> | | | | | | | |
| <i>Setaria parviflora</i> | | | | | | | |
| <i>Sida acuta</i> | | | | | | | |
| <i>Sida ciliaris</i> | | | | | | | |
| <i>Sideroxylon salicifolium</i> | | | | | | | |
| <i>Solanum elaeagnifolium</i> | | | | | | | |
| <i>Solidago gigantea</i> | | | | | | | |
| <i>Solidago sempervirens</i> | | | | | | | |
| <i>Solidago stricta</i> | | | | | | | |
| <i>Sorghum halepense</i> | No | Mesic | <1 | 1-100 | Throughout | Mediterranean region | BW 307 |
| <i>Spartina bakeri</i> | | | | | | | |
| <i>Spermacoce assurgens</i> | | | | | | | |
| <i>Spermacoce verticillata</i> | No | Mesic | 1-100 | 10,001-100,000 | South, Central, North | Africa | |
| <i>Sporobolus indicus</i> var. <i>pyramidalis</i> | No | Mesic | 101-1,000 | 10,001-100,000 | South, Central, Panhandle | Tropical America | |
| <i>Stachytarpheta jamaicensis</i> | | | | | | | |
| <i>Stenotaphrum secundatum</i> | | | | | | | |
| <i>Stylosanthes hamata</i> | | | | | | | |
| <i>Swertia mahagoni</i> | No | Mesic | <1 | 1-100 | South | Extreme Southern Florida to South America | |
| <i>Syagrus romanzoffianum</i> | | | | | | | |
| <i>Syzygium cumini</i> | No | Mesic | 1-100 | 1,001-10,000 | South | Asia | |
| <i>Taxodium ascendens</i> | | | | | | | |
| <i>Taxodium distichum</i> | | | | | | | |
| <i>Teucrium canadense</i> | | | | | | | |
| <i>Thalia geniculata</i> | | | | | | | |
| <i>Thelypteris interrupia</i> | | | | | | | |
| <i>Thelypteris kunthii</i> | | | | | | | |

Water Conservation Area 2 - Preliminary List of Vascular Plants

Bold = Exotic, *Ifalics* = Cultivated

| | Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|------------|---------|-----------|----------------|---------------------------|------------------|---------|
| <i>Thelypteris palustris</i> var. <i>pubescens</i> | | | | | | | |
| <i>Tillandsia recurvata</i> | | | | | | | |
| <i>Trema micranthum</i> | | | | | | | |
| <i>Tribulus cistoides</i> | No | Mesic | <1 | 1-100 | South, Central | Tropical America | |
| <i>Tridax procumbens</i> | No | Mesic | <1 | 1,001-10,000 | South, Central | Tropical America | BW 303 |
| <i>Typha domingensis</i> | | | | | | | |
| <i>Urena lobata</i> | | | | | | | |
| <i>Urochloa mutica</i> | Yes | Wet | 101-1,000 | 10,001-100,000 | South, Central | Africa | |
| <i>Utricularia foliosa</i> | | | | | | | |
| <i>Vallisneria americana</i> | | | | | | | |
| <i>Verbena scabra</i> | | | | | | | |
| <i>Vitis cinerea</i> var. <i>floridana</i> | | | | | | | |
| <i>Vitis rotundifolia</i> | | | | | | | |
| <i>Vitis shuttleworthii</i> | | | | | | | |
| <i>Waltheria indica</i> | | | | | | | |
| <i>Wedelia trilobata</i> | No | Mesic | <1 | 101-1,000 | South, Central, Panhandle | West Indies | |
| <i>Youngia japonica</i> | No | Mesic | <1 | >100,000 | Throughout | SE Asia | BW 279 |
| <i>Zizaniopsis miliacea</i> | | | | | | | |

Water Conservation Area 2 - Preliminary List of Vascular Plants

Bold = Exotic, *Italics* = Cultivated

| Nat. Area? | Habitat | Acreage | Pop. Size | Florida Range | Natural Range | Voucher |
|--|---------|---------|-----------|----------------|---------------------------|---------|
| <i>Thelypteris palustris</i> var. <i>pubescens</i> | | | | | | |
| <i>Tillandsia recurvata</i> | | | | | | |
| <i>Trema micranthum</i> | | | | | | |
| <i>Tribulus cistoides</i> | No | Mesic | <1 | 1-100 | South, Central | BW 303 |
| <i>Tridax procumbens</i> | No | Mesic | <1 | 1,001-10,000 | South, Central | |
| <i>Typha domingensis</i> | | | | | Tropical America | |
| <i>Urena lobata</i> | | | | | Tropical America | |
| <i>Urochloa mutica</i> | Yes | Wet | 101-1,000 | 10,001-100,000 | South, Central | |
| <i>Utricularia foliosa</i> | | | | | Africa | |
| <i>Vallisneria americana</i> | | | | | | |
| <i>Verbena scabra</i> | | | | | | |
| <i>Vitis cinerea</i> var. <i>floridana</i> | | | | | | |
| <i>Vitis rotundifolia</i> | | | | | | |
| <i>Vitis shuttleworthii</i> | | | | | | |
| <i>Waltheria indica</i> | | | | | | |
| <i>Wedelia trilobata</i> | No | Mesic | <1 | 101-1,000 | South, Central, Panhandle | BW 279 |
| <i>Youngia Japonica</i> | No | Mesic | <1 | >10,000 | Throughout | |
| <i>Zizaniopsis miliacea</i> | | | | | SE Asia | |

Table I - Exotic Plant Taxa

| Taxon Name | # Sites |
|---|---------|
| <i>Abrus precatorius</i> | 1 |
| <i>Acacia auriculiformis</i> | 3 |
| <i>Acalypha alopecuroides</i> | 1 |
| <i>Ageratum houstonianum</i> | 1 |
| <i>Albizia lebbeck</i> | 2 |
| <i>Alternanthera paronychoides</i> | 1 |
| <i>Alternanthera philoxeroides</i> | 5 |
| <i>Alternanthera sessilis</i> | 1 |
| <i>Alysicarpus ovalifolius</i> | 1 |
| <i>Amaranthus spinosus</i> | 3 |
| <i>Amaranthus viridis</i> | 3 |
| <i>Anagallis arvensis</i> | 1 |
| <i>Antigonon leptopus</i> | 1 |
| <i>Ardisia elliptica</i> | 1 |
| <i>Asclepias curassavica</i> | 2 |
| <i>Bischofia javanica</i> | 5 |
| <i>Blechum pyramidatum</i> | 3 |
| <i>Bothriochloa pertusa</i> | 1 |
| <i>Brassica juncea</i> | 1 |
| <i>Bulbostylis barbata</i> | 1 |
| <i>Callisia fragrans</i> | 1 |
| <i>Caperonia palustris</i> | 1 |
| <i>Cardiospermum halicacabum</i> | 2 |
| <i>Carica papaya</i> | 4 |
| <i>Caryota mitis</i> | 2 |
| <i>Casuarina equisetifolia</i> | 4 |
| <i>Casuarina glauca</i> | 2 |
| <i>Catharanthus roseus</i> | 2 |
| <i>Cestrum diurnum</i> | 2 |
| <i>Chamaesyce lasiocarpa</i> | 1 |
| <i>Chenopodium album</i> | 1 |
| <i>Chenopodium ambrosioides</i> | 2 |
| <i>Chenopodium murale</i> | 1 |
| <i>Chloris cf. radiata</i> | 1 |
| <i>Citrus aurantium</i> | 1 |
| <i>Citrus sinensis</i> | 1 |
| <i>Citrus x paradisi</i> | 2 |
| <i>Colocasia esculenta</i> | 4 |
| <i>Commelinia caroliniana</i> | 1 |
| <i>Conobea multifida</i> | 1 |
| <i>Corchorus aestuans</i> | 2 |
| <i>Cordia myxa</i> | 1 |
| <i>Coronopus didymus</i> | 1 |
| <i>Crinum sp.</i> | 1 |
| <i>Crotalaria lanceolata</i> | 3 |
| <i>Crotalaria pallida</i> var. <i>obovata</i> | 3 |
| <i>Crotalaria spectabilis</i> | 2 |
| <i>Cucumis melo</i> | 2 |
| <i>Cupaniopsis anacardioides</i> | 3 |
| <i>Cuphea carthagenensis</i> | 4 |
| <i>Cynodon dactylon</i> | 6 |
| <i>Cyperus esculentus</i> | 3 |
| <i>Cyperus iria</i> | 1 |
| <i>Cyperus ochraceus</i> | 1 |
| <i>Cyperus prolifer</i> | 1 |
| <i>Cyperus rotundus</i> | 2 |
| <i>Cyperus sphacelatus</i> | 1 |
| <i>Dactyloctenium aegyptium</i> | 6 |
| <i>Delonix regia</i> | 1 |
| <i>Desmanthus cf. leptophyllus</i> | 1 |
| <i>Desmodium triflorum</i> | 4 |
| <i>Echinochloa colona</i> | 2 |
| <i>Echinochloa crusgalli</i> | 1 |
| <i>Eichornia crassipes</i> | 4 |
| <i>Eleocharis nigrescens</i> | 2 |
| <i>Eleusine indica</i> | 4 |
| <i>Emilia fosbergii</i> | 3 |
| <i>Emilia sonchifolia</i> | 2 |
| <i>Epipremnum pinnatum</i> | 1 |

Table I - Exotic Plant Taxa

| | |
|---|---|
| <i>Pteris vittata</i> | 6 |
| <i>Pueraria montana</i> var. <i>lobata</i> | 1 |
| <i>Rhynchelytrum repens</i> | 5 |
| <i>Richardia brasiliensis</i> | 1 |
| <i>Richardia grandiflora</i> | 1 |
| <i>Richardia scabra</i> | 1 |
| <i>Ricinus communis</i> | 3 |
| <i>Rottboellia cochinchinensis</i> | 2 |
| <i>Roystonea regia</i> | 2 |
| <i>Sacciolepis indica</i> | 3 |
| <i>Sansevieria hyacinthoides</i> | 1 |
| <i>Schefflera actinophylla</i> | 2 |
| <i>Schinus terebinthifolius</i> | 7 |
| <i>Senna alata</i> | 1 |
| <i>Senna occidentalis</i> | 1 |
| <i>Senna pendula</i> var. <i>glabrata</i> | 4 |
| <i>Sida cordifolia</i> | 2 |
| <i>Solanum cf. torvum</i> | 1 |
| <i>Solanum diphyllum</i> | 1 |
| <i>Solanum tampicense</i> | 1 |
| <i>Solanum viarum</i> | 2 |
| <i>Sonchus asper</i> | 2 |
| <i>Sonchus oleraceus</i> | 2 |
| <i>Sorghum arundinaceum</i> | 1 |
| <i>Sorghum bicolor</i> | 1 |
| <i>Sorghum halepense</i> | 3 |
| <i>Spermacoce verticillata</i> | 5 |
| <i>Spirodela punctata</i> | 3 |
| <i>Sporobolus indicus</i> var. <i>indicus</i> | 5 |
| <i>Sporobolus indicus</i> var. <i>pyramidalis</i> | 5 |
| <i>Swietenia mahagoni</i> | 1 |
| <i>Syagrus romanzoffianum</i> | 1 |
| <i>Syngonium podophyllum</i> | 1 |
| <i>Syzygium cumini</i> | 3 |
| <i>Syzygium jambos</i> | 1 |
| <i>Thelypteris dentata</i> | 2 |
| <i>Tradescantia pallida</i> | 1 |
| <i>Tribulus cistoides</i> | 1 |
| <i>Tridax procumbens</i> | 3 |
| <i>Urochloa mutica</i> | 5 |
| <i>Urochloa plantaginea</i> | 1 |
| <i>Urochloa ramosa</i> | 1 |
| <i>Urochloa subquadripala</i> | 1 |
| <i>Verbena bonariensis</i> | 2 |
| <i>Veronica cinerea</i> | 2 |
| <i>Washingtonia robusta</i> | 2 |
| <i>Wedelia trilobata</i> | 4 |
| <i>Xyris jupicai</i> | 4 |
| <i>Younghia japonica</i> | 4 |

Table III - Endangered and Threatened Taxa

| Taxon Name | # Sites | Status |
|--|---------|--------|
| <i>Anacheilium cochleatum</i> var. <i>triandrum</i> | 1 | E |
| <i>Anemia wrightii</i> | 1 | E |
| <i>Bletia purpurea</i> | 1 | T |
| <i>Byrsonima lucida</i> | 1 | E |
| <i>Campyloneurum phyllitidis</i> | 2 | E |
| <i>Chrysophyllum oliviforme</i> | 1 | E |
| <i>Cucurbita okeechobeensis</i> | 1 | E |
| <i>Epidendrum rigidum</i> | 1 | E |
| <i>Eulophia alta</i> | 5 | T |
| <i>Harrisella filiformis</i> | 1 | T |
| <i>Ilex krugiana</i> | 1 | E |
| <i>Jacquemontia curtissii</i> | 1 | E |
| <i>Lechea cernua</i> | 1 | T |
| <i>Lilium catesbaei</i> | 3 | T |
| <i>Melanthera parvifolia</i> | 1 | E |
| <i>Nemastylis floridana</i> | 1 | E |
| <i>Odontosoria clavata</i> | 1 | E |
| <i>Ophioglossum palmatum</i> | 1 | E |
| <i>Platanthera nivea</i> | 1 | T |
| <i>Polygala smallii</i> | 1 | E |
| <i>Polystachya concreta</i> | 1 | E |
| <i>Selaginella eatonii</i> | 1 | E |
| <i>Thelypteris serrata</i> | 1 | E |
| <i>Tillandsia balbisiana</i> | 5 | T |
| <i>Tillandsia fasciculata</i> var. <i>densispica</i> | 4 | E |
| <i>Tillandsia flexuosa</i> | 1 | E |
| <i>Tillandsia utriculata</i> | 6 | E |
| <i>Tillandsia variabilis</i> | 2 | T |

Table II - Invasive Taxa

| Taxon name | # Sites Where Invasive | Tot. # Sites | % Sites Where Invasive |
|---|------------------------|--------------|------------------------|
| <i>Abrus precatorius</i> | 1 | 1 | 100% |
| <i>Alternanthera philoxeroides</i> | 5 | 5 | 100% |
| <i>Alternanthera sessiliis</i> | 1 | 1 | 100% |
| <i>Bischofia javanica</i> | 2 | 5 | 40% |
| <i>Carica papaya</i> | 4 | 4 | 100% |
| <i>Casuarina equisetifolia</i> | 1 | 4 | 25% |
| <i>Citrus aurantium</i> | 1 | 1 | 100% |
| <i>Citrus sinensis</i> | 1 | 1 | 100% |
| <i>Citrus x paradisi</i> | 1 | 2 | 50% |
| <i>Colocasia esculenta</i> | 2 | 4 | 50% |
| <i>Cordia myxa</i> | 1 | 1 | 100% |
| <i>Cupaniopsis anacardioides</i> | 1 | 3 | 33% |
| <i>Cuphea carthagenensis</i> | 2 | 4 | 50% |
| <i>Eichhornia crassipes</i> | 4 | 4 | 100% |
| <i>Eleocharis nigrescens</i> | 2 | 2 | 100% |
| <i>Epipremnum pinnatum</i> | 1 | 1 | 100% |
| <i>Eragrostis atrovirens</i> | 3 | 3 | 100% |
| <i>Eugenia uniflora</i> | 2 | 3 | 67% |
| <i>Ficus microcarpa</i> | 3 | 4 | 75% |
| <i>Hydrilla verticillata</i> | 4 | 4 | 100% |
| <i>Hymenachne amplexicaulis</i> | 2 | 2 | 100% |
| <i>Hyptis verticillata</i> | 1 | 3 | 33% |
| <i>Lantana camara</i> | 1 | 4 | 25% |
| <i>Limnophila sessiliflora</i> | 1 | 1 | 100% |
| <i>Lipocarpha aristulata</i> | 1 | 1 | 100% |
| <i>Lygodium microphyllum</i> | 3 | 3 | 100% |
| <i>Mangifera indica</i> | 1 | 1 | 100% |
| <i>Melaleuca quinquenervia</i> | 6 | 6 | 100% |
| <i>Mimosa pigra</i> | 1 | 1 | 100% |
| <i>Momordica charantia</i> | 1 | 5 | 20% |
| <i>Najas cf. minor</i> | 1 | 1 | 100% |
| <i>Nephrolepis cordifolia</i> | 3 | 3 | 100% |
| <i>Neyraudia reynaudiana</i> | 1 | 2 | 50% |
| <i>Oeceoclades maculata</i> | 1 | 1 | 100% |
| <i>Panicum repens</i> | 7 | 7 | 100% |
| <i>Paspalum notatum</i> | 1 | 4 | 25% |
| <i>Passiflora edulis</i> | 1 | 1 | 100% |
| <i>Pistia stratiotes</i> | 5 | 5 | 100% |
| <i>Pouzolzia zeylanica</i> | 1 | 1 | 100% |
| <i>Psidium cattleianum</i> | 2 | 2 | 100% |
| <i>Psidium guajava</i> | 4 | 4 | 100% |
| <i>Pteris vittata</i> | 4 | 6 | 67% |
| <i>Ricinus communis</i> | 2 | 3 | 67% |
| <i>Roystonea regia</i> | 1 | 2 | 50% |
| <i>Sacciolepis indica</i> | 3 | 3 | 100% |
| <i>Schefflera actinophylla</i> | 1 | 2 | 50% |
| <i>Schinus terebinthifolius</i> | 7 | 7 | 100% |
| <i>Senna pendula</i> var. <i>glabrata</i> | 2 | 4 | 50% |
| <i>Solanum cf. torvum</i> | 1 | 1 | 100% |
| <i>Solanum tampicense</i> | 1 | 1 | 100% |
| <i>Solanum viarum</i> | 1 | 2 | 50% |
| <i>Spermacoce verticillata</i> | 2 | 5 | 40% |
| <i>Spirodela punctata</i> | 2 | 3 | 67% |
| <i>Syngonium podophyllum</i> | 1 | 1 | 100% |
| <i>Syzygium cumini</i> | 1 | 3 | 33% |
| <i>Thelypteris dentata</i> | 2 | 2 | 100% |
| <i>Tridax procumbens</i> | 1 | 3 | 33% |
| <i>Urochloa mutica</i> | 4 | 5 | 80% |
| <i>Vernonia cinerea</i> | 1 | 2 | 50% |
| <i>Xyris jupicai</i> | 4 | 4 | 100% |
| <i>Youngia japonica</i> | 2 | 4 | 50% |